

UTMD-049

"Near" weighted utilitarian characterizations of

Pareto optima

Yeon-Koo Che Columbia University

Jinwoo Kim Seoul National University

> Fuhito Kojima University of Tokyo

Christopher Thomas Ryan University of British Columbia

March 28, 2023

"Near" weighted utilitarian characterizations of Pareto optima¹

Yeon-Koo Che² Jinwoo Kim³ Fuhito Kojima⁴ Christopher Thomas Ryan⁵

March 28, 2023

Abstract

We characterize Pareto optimality via "near" weighted utilitarian welfare maximization. One characterization sequentially maximizes utilitarian welfare functions using a finite sequence of nonnegative and eventually positive welfare weights. The other maximizes a utilitarian welfare function with a certain class of positive hyperreal weights. The social welfare ordering represented by these "near" weighted utilitarian welfare criteria are characterized by the standard axioms for weighted utilitarianism under a suitable weakening of the continuity axiom.

Keywords: Pareto optima, weighted utilitarian welfare maximization, sequential utilitarian welfare maximization, simple hyperreal weights, weak continuity

JEL Numbers: C60, D60, D50.

- ³Department of Economics and SIER, Seoul National University
- ⁴Department of Economics and Market Design Center, University of Tokyo
- ⁵Operations and Logistics Division, UBC Sauder School of Business, University of British Columbia

¹We thank the Co-editor, Asher Wolinsky, as well as four anonymous referees for comments that led to major improvements. We are also grateful to Florian Brandl, Timothy Y Chan, Alexander Engau, Atsushi Kajii, Yuichiro Kamada, Michihiro Kandori, and Eitetsu Ken for their helpful comments and conversations, as well as to seminar audiences at Arizona State, Carlo Alberto, Carlos III, Carnegie Mellon, Harvard-MIT, KAIST, Northwestern, Stanford, University of British Columbia, UC-Davis, and the International Conference on Game Theory at Stony Brook. We are especially grateful to Ludvig Sinander and Gregorio Curello for a question that led to this project. We acknowledge research assistance from Yutaro Akita, Nanami Aoi, Xuandong Chen, William Grimme, Jiangze Han, Yusuke Iwase, Masanori Kobayashi, Kevin Li, Leo Nonaka, Ryo Shirakawa, Shoya Tsuruta, Ayano Yago, and Yutong Zhang. Yeon-Koo Che is supported by National Science Foundation Grant SES-1851821. Fuhito Kojima is supported by the JSPS KAKENHI Grant-In-Aid 21H04979. Christopher Thomas Ryan is supported by the Natural Sciences and Engineering Research Council of Canada Discovery Grant RGPIN-2020-06488 and the UBC Sauder Exploratory Grants Program. This work was supported by the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2020S1A5A2A03043516).

²Department of Economics, Columbia University

1 Introduction

Pareto optimality is a central concept in economics for its normative appeal. Also central is weighted utilitarian welfare maximization; e.g., Harsanyi (1955) famously defended it as a social welfare function based on several normative axioms. Moreover, weighted utilitarianism is widely invoked in practice, including applied research and policy debates. Given the prominent roles played by these two concepts, attempts have been made to establish a connection between Pareto optima and weighted utilitarianism—or more precisely, a characterization of Pareto optima via weighted utilitarian welfare maximization. Yet, such a characterization has so far been elusive.

It is well known that, given a closed and convex utility possibility set, which we assume throughout, every Pareto optimal utility vector maximizes some *nonnegatively* weighted sum of utilities of agents (see Proposition 3.45 in Bewley (2009)). But the converse is false: not every such maximizer is Pareto optimal. To see this, suppose a society consists of two agents, 1 and 2, and the utility possibility set is given by U in Figure 1. All points on



Figure 1: Weighted utilitarian welfare maximization need not yield a Pareto optimum.

the "outer" boundary, including the vertical segment, maximize suitably weighted sums of agents' utilities within U, but not all of them are Pareto optimal. In particular, the points on the vertical segment strictly below u, such as u''', all maximize the utility sum with weights $\phi = (1, 0)$ —i.e., only 1's utility. Yet, none of these points is Pareto optimal. The reason is that the welfare of the agent receiving zero weight is *not* counted.

By contrast, if weights are restricted to be (strictly) positive for all agents, weighted utilitarian welfare maximization does always yield a Pareto optimum (Proposition 3.23 of Bewley (2009)). But the converse is false: not every Pareto optimal outcome can be obtained in this way. In Figure 1, u' is Pareto optimal and obtained by weighted utilitarian welfare maximization with positive weights, but u and u'', which are also Pareto optimal, cannot be obtained.

While positive welfare weights do not yield points like u in Figure 1, one may conjecture that they may "in the limit"; for instance, u is a limit of welfare-maximizing utility vectors with positive weights (1, 1/n), as $n \to \infty$. Indeed, Arrow, Barankin, and Blackwell (1953) show that every Pareto optimal vector is a limit of a sequence of utility vectors that



Figure 2: The "tilted cone" adapted from Arrow, Barankin, and Blackwell (1953) and Bitran and Magnanti (1979). The set is the convex hull of the portion of the unit disk centered at the origin in the u_1 - u_2 plane from point K to point S (where $\alpha^2 + \beta^2 = 1$ with $\alpha \in (0, 1)$) and the apex point V = (0, 1, 1). The blue surface, including all of its boundaries except for the dotted line, is the set of Pareto optimal utility vectors.

maximize some positively weighted sum of utilities—a result known as the ABB theorem.¹ Unfortunately, this too does not lead to a characterization when there are more than two agents:² again its converse is false—namely, a limit point of such a sequence may not be Pareto optimal. To see this, suppose there are three agents, 1, 2, and 3, with possible utility vectors depicted in Figure 2. The point K is a limit of the sequence of points maximizing a positively weighted sum of utilities (see the arrow) but is Pareto dominated, say, by the point V. The relationship between Pareto optima and the alternative notions of weighted utilitarianism is depicted in Figure 3, where U^P is the set of Pareto optimal utility vectors while U^+ and U^{++} are the sets of utility vectors that maximize nonnegatively weighted and (strict) positively weighted utilitarian welfare, respectively, with $cl(U^{++})$ being the closure of U^{++} .

This paper provides exact characterizations of Pareto optima by close variants of weighted

¹This theorem has spawned a series of extensions to spaces more general than Euclidean space. See Daniilidis (2000) for a survey of ABB theorems.

²When there are two agents, the limit $u \in U$ of any sequence $\{u^k\}$ of utilities $u^k \in U$ maximizing a positively weighted sum of utilities is Pareto optimal, where U is the utility possibility set, assumed to be closed and convex. To see it, let $\{\phi^k\}$ be the sequence of positive weights, normalized to be in the simplex, such that $u^k \in \arg\max_{(u'_1, u'_2) \in U} \sum_{i=1}^2 \phi_i^k u'_i$, and let ϕ denote its limit (say of a convergent subsequence). Clearly, $u \in \arg\max_{(u'_1, u'_2) \in U} \sum_{i=1}^2 \phi_i u'_i$. If ϕ_1 and ϕ_2 are both positive, then u is Pareto optimal, so assume $\phi_1 = 1$ and $\phi_2 = 0$ without loss. Suppose for contradiction u is not Pareto optimal. Then, there must exist $v \in U$ such that $v_1 = u_1$ and $v_2 > u_2$, where the equality holds since $u \in \arg\max_{(u'_1, u'_2) \in U} \sum_{i=1}^2 \phi_i u'_i = \arg\max_{(u'_1, u'_2) \in U} u'_1$. Since u^k 's are all Pareto optimal, we have $u_1^k \leq v_1 = u_1$ and $u_2^k \geq v_2 > u_2$ for all k, so u^k never converges to u, a contradiction.



Figure 3: Alternative notions of utilitarian welfare maximization in relationship with Pareto optimality. The containment $U^{++} \subset U^P \subset U^+$ follows from Propositions 3.23 and 3.45 in Bewley (2009). The containment $U^P \subset cl(U^{++})$ is from Arrow, Barankin, and Blackwell (1953). The containment $cl(U^{++}) \subset U^+$ is straightforward.

utilitarian welfare maximization. To ease language, we will refer to weighted utilitarian welfare maximization simply as utilitarianism.³

We show that a utility vector u is Pareto optimal if and only if there exists a finite sequence of nonnegative and "eventually positive" welfare weights such that in each round t, u maximizes the round-t weighted sum of utilities out of those surviving from round t-1. Here, "eventually positive" means that the support of the weight vector strictly grows over the rounds with the weight vector in the final round having full support.

To illustrate why our SUWM successfully characterizes Pareto optima, let us revisit why neither the "nonnegative utilitarianism" captured by U^+ nor the "positive utilitarianism" captured by U^{++} in Figure 3 works. Nonnegative utilitarianism can include Pareto suboptimal outcomes because some individual's utility may not "count" at all. Positive utilitarianism avoids this problem by requiring that every individual's utility carry positive weights. However, as can be seen from Figure 1, it excludes Pareto optimal outcomes that can be achieved only by assigning some individuals "infinitely smaller" weights than others. SUWM resolves this seeming conflict by assigning positive weights to individuals so that "every agent's welfare counts" but in different rounds: Individuals with strictly positive weights only in later rounds can be regarded as carrying infinitely smaller weights than those with positive weights in earlier rounds.

The preceding observation gives rise to our second characterization of Pareto optimality, via one-shot maximization of utilitarian welfare with *hyperreal* weights. Hyperreal numbers

³In particular, note that we drop the qualifier "weighted" in our usage of the concept of utilitarianism while keeping in mind that utilitarianism is always used in the weighted sense. Indeed, we have no occasion to discuss unweighted utilitarianism. It is only for emphasis or to provide further clarification that we use the qualifier "weighted" in connection to utilitarianism.

include not only standard real numbers but also infinitesimal "numbers." The space of hyperreal numbers is very large, which may limit the usefulness of the characterization. By contrast, our characterization places an added discipline and structure on such social welfare functions. The resulting criterion, called *simple hyperreal utilitarian welfare maximization* (SHUWM), requires the hyperreal weights to be not only strictly positive but also represented by a finite sequence of nonnegative and eventually positive real weights.

Both of these characterizations of Pareto optimality capture the essential feature of standard weighted utilitarian welfare maximization. First, SUWM and SHUWM reduce to utilitarianism in many situations in which the former involves one-round maximization and the latter involves no infinitesimal weights. Second, the welfare functions used in these characterizations are inherently *linear* (based on weighted sums of agent utilities), albeit with SUWM having several rounds of linear optimizations and SHUWM involving hyperreal weights. Third, a consequence of this linearity is that the utilities of individuals are aggregated by weights that do not depend on the particular utility profile under consideration, a property we refer to as having "constant weights." This is in contrast to other social welfare functions, such as Rawlsian and leximin whose weighting of an agent's utility depends on her relative position in a given utility vector.

The sense in which our characterizations constitute "near" utilitarianism is further clarified by the social welfare orderings that underpin our characterizations. d'Aspremont and Gevers (2002) show that for social welfare orderings to be represented by a utilitarian welfare function, they must not only satisfy the **Pareto Principle**—namely, they must preserve Pareto domination order—but they must also satisfy two additional axioms: **Invariance** and **Continuity**. **Invariance** requires the orderings to be robust to translation and/or scaling of the utility profiles of individuals. **Continuity** requires the orderings to be robust to perturbations of utility profiles. **Continuity** effectively forces the welfare weights of agents to be in the same order of magnitude, thus making it impossible for the weight of an agent to be infinitesimally smaller than that of another agent. Since the latter feature is crucial for characterizing Pareto optima, **Continuity** must be relaxed.

Indeed, we show that the welfare orderings associated with SUWM and SHUWM can be obtained by the same set of axioms under a suitable weakening of Continuity—more precisely, by the Pareto Principle, Invariance, and Weak Continuity. The last axiom weakens Continuity by requiring welfare orderings to be robust to perturbations of utilities of *some*, *but not necessarily all*, individuals, which is in line with our characterization of Pareto optima that allows some individuals to be assigned infinitely larger weights than others. That our welfare notions preserve a version of continuity, albeit weakened, is a nontrivial marker of the sense in which SUWM and SHUWM closely resemble utilitarianism. In particular, the same marker is not shared with other reasonable characterizations. For instance, as we show in a subsequent section, an (unrestricted) hyperreal-weighted utilitarian welfare function does not satisfy Weak Continuity.

Our characterizations of Pareto optimality fulfill a long-standing intellectual pursuit of providing a weighted utilitarian foundation for Pareto optimality. In addition, our characterizations of Pareto optimality serve other useful purposes. First, the SUWM characterization could provide a tractable method for computing Pareto optimal allocations, which may be useful in the market design context. In fact, SUWM can be viewed as a generalization of the serial dictatorship mechanism in which each agent acts sequentially according to serial order to maximize her utility. Serial dictatorship is used widely for Pareto optimally allocating indivisible resources when monetary transfers cannot be used. For instance, serial dictatorship with a randomized serial order—known as random serial dictatorship—is used for assigning public school seats, public and campus housing, and human organs. One could imagine that SUWM can serve a similar practical purpose, but in a much more general setting that goes beyond a one-to-one assignment. In each round, one can let a group of agents negotiate over feasible allocations at that round, as will be made precise in Section 5. Indeed, a procedure like this is used in the assignment of campus housing.⁴ Alternatively, a central clearinghouse may compute an optimal choice for the group in each round.⁵

Second, the SUWM characterization could serve as a useful analytical tool for analyzing the behavior of Pareto optima as a set. For instance, one may study the comparative statics of Pareto optima—i.e., how they change as the primitives change—utilizing monotone comparative statics methods developed for optimization (e.g., Topkis (1998) and Milgrom and Shannon (1994)). The "round-wise" linear structure of SUWM admits a convenient aggregation property that is crucial for such an analysis. Indeed, Che, Kim, and Kojima (2019) use this property to develop a theory of monotone comparative statics of Pareto optima: they show that when agents' utility functions shift in a way that leads to higher *individual* choices of decisions (e.g., Milgrom and Shannon (1994)), the Pareto optima shift to a higher set of actions in a suitable sense.⁶

The remainder of the paper is organized as follows. Section 2 describes our setting and establishes a few preliminaries used in our main results. Section 3 states our characterization of Pareto optimality. Here, we discuss the tools used to prove the result. Section 4 establishes the axiomatization of SUWM and SHUWM. Section 5 looks at other reasonable characterizations of Pareto optimality that fail at least one of the "near" utilitarian axioms set out in the previous section. Section 6 concludes with some suggestions for future work. The appendix provides proofs of our main characterization and axiomatization results. A supplementary appendix contains statements and proofs of additional results.

⁴ For example, the campus housing assignment at Columbia university uses a cohort-based serial dictatorship, in which a group of students chooses a suite collectively in each round of the serial dictatorship procedure; presumably, the students then negotiate among themselves to allocate rooms within the assigned suite.

⁵In both scenarios, we are implicitly assuming complete information. In case agents' preferences are unobserved, the designer must rely on their preference reports, in which case agents' incentives become an important aspect of the market design. While this issue is beyond the scope of the current paper, it can be addressed in some specific settings such as cohort-based serial dictatorship mentioned in Footnote 4, where the standard strategy-proofness property would extend to a group of students as long as they know their preferences.

⁶In particular, properties such as supermodularity and increasing differences, which are important for the monotone comparative statics analysis, are preserved under this aggregation. The same proof would not have been possible with nonlinear welfare functions.

2 Setting and preliminaries

In this section, we introduce our basic setting and introduce some elementary concepts needed for stating our main results.

Let $I = \{1, 2, ..., n\}$ denote a finite set of agents and the *utility possibility set* $U \subset \mathbb{R}^n$ be the set of possible utility vectors the agents may attain. We assume that U is closed and convex. If U stems from an underlying choice space X via utility functions $(u_i)_{i \in I} : X \to \mathbb{R}^n$, then we let

$$U = \{ u \in \mathbb{R}^n \mid u \le u(x) \text{ for some } x \in X \}.^7$$
(1)

That U is closed and convex is arguably a mild assumption that is satisfied if, for instance, U is induced by utility functions $(u_i)_{i \in I}$ that are upper semicontinuous and concave on a choice set X that is compact and convex.⁸

For any $u, v \in \mathbb{R}^n$, we write $v \ge u$ if $v_i \ge u_i$ for all $i \in I$, v > u if $v \ge u$ and $v \ne u$, and $v \gg u$ if $v_i > u_i$ for all $i \in I$. We say a point u in U is *Pareto optimal* with respect to U if there exists no $v \in U$ with v > u. Let $U^P \subset U$ denote the set of all Pareto optimal points (or, more simply, Pareto optima).

For any $\phi \in \mathbb{R}^n$, consider the optimization problem:

$$\max_{u \in U} \langle \phi, u \rangle, \tag{2}$$

where $\langle \phi, u \rangle := \sum_{i=1}^{n} \phi_i u_i$. We call ϕ a weight vector. Throughout the paper, we only consider nonzero weight vectors (i.e., $\phi \neq 0$). We say a point $u \in U$ maximizes the weight vector ϕ over U (or simply maximizes ϕ) if u is a solution to (2). We call a weight vector ϕ nonnegative if $\phi > 0$ and positive if $\phi \gg 0$. For any vector $v \in \mathbb{R}^n$, the support of v is the set of indices where v is nonzero; i.e., $\operatorname{supp} v := \{i \in I \mid v_i \neq 0\}$. A positive ϕ has full support; i.e., $\operatorname{supp} \phi = I$.

Our discussion uses the language of hyperreal numbers. We introduce the basics here. The set of hyperreal numbers $*\mathbb{R}$ consists of real numbers as well as "infinite" and "infinitesimal" numbers. Infinite numbers are larger than any real number. Infinitesimal numbers (or simply infinitesimals) are closer to 0 than any real number. A formal definition of $*\mathbb{R}$ is somewhat tedious and we will not reproduce it here. Instead, we refer the reader to Goldblatt (2012). Although the use of hyperreal numbers (in what is termed *nonstandard analysis*) is not completely standard in economics, it has been used in a variety of settings including choice under uncertainty (Blume, Brandenburger, and Dekel, 1991), game theory (Dilmé, 2022), and exchange economies (Brown and Robinson, 1975). See Anderson (1991) for a survey of applications of nonstandard analysis to economics.

Important properties of the set of hyperreal numbers for our purposes are that (i) \mathbb{R} contains a (positive) infinitesimal number, i.e., an element $\epsilon \in \mathbb{R}$ such that $\epsilon < r$ for every

⁷To be precise, the utility possibility set is often defined as $\{u \in \mathbb{R}^n \mid u = (u_i(x))_{i \in I} \text{ for some } x \in X\}$, which differs from (1). However, the two sets share the same set of Pareto optima since those points are on the common outer boundary of the sets. Thus, formulating the set U either way makes no difference for our results while the current formation facilitates our analysis.

⁸Note that compactness and convexity of the choice set X are satisfied if, for instance, all lotteries of social outcomes, which are in turn finite, or more generally compact, are feasible.

positive real number r while $\epsilon > 0$, and that (ii) arithmetic operations such as addition and multiplication, as well as order relations, are well defined and extended from \mathbb{R} to $*\mathbb{R}$ in expected ways.

3 Characterizations of Pareto optimality

This section presents our main result, Theorem 1, that provides two alternative "near" (weighted) utilitarian characterizations of the set U^P of Pareto optimal points of a given closed convex set U. To state these characterizations, we first introduce some additional terminology and definitions. These definitions will be interpreted after the statement of Theorem 1.

A sequence $\Phi = (\phi^1, \phi^2, \dots, \phi^T)$ of weight vectors is *nonnegative* if ϕ^t is nonnegative for every $t \in \{1, \dots, T\}$. We say that a sequence Φ of weight vectors is *eventually positive* if $\operatorname{supp} \phi^{t-1} \subsetneq \operatorname{supp} \phi^t$ for all $t = 2, \dots, T$ and $\operatorname{supp} \phi^T = I$. Note that eventual positivity implies $T \leq n$ since the support *strictly* grows along the sequence.

Definition 1 (Sequential utilitarian welfare maximization (SUWM)). We say $u \in U$ sequentially maximizes a sequence $\Phi = (\phi^1, \phi^2, \dots, \phi^T)$ of weight vectors over U if

$$u \in U^t := \arg \max_{u' \in U^{t-1}} \langle \phi^t, u' \rangle$$
, for each $t = 1, \dots, T$, (3)

where $U^0 = U$. We say $u \in U$ sequentially maximizes utilitarian welfare over U—or, more simply, u is an SUWM solution of U—if there exists a sequence Φ of nonnegative and eventually positive weight vectors such that u sequentially maximizes Φ .

The following definition uses the concept of hyperreals introduced in the preliminaries section. We call a vector $\phi \in ({}^*\mathbb{R})^n$ of hyperreal weights *simple* if there exists a positive infinitesimal number ϵ and a sequence $\Phi = (\phi^1, \phi^2, \dots, \phi^T)$ of nonnegative and eventually positive weight vectors in \mathbb{R}^n such that

$$\phi = \sum_{t \in \{1,\dots,T\}} \epsilon^{t-1} \phi^t.$$
(4)

An example with two individuals can illustrate the restriction associated with a "simple" hyperreal vector. Consider the hyperreal weight vector $(1 + \epsilon, 1)$, with ϵ being a positive infinitesimal number. This vector is not simple. To see this, note that the only way to express the vector $(1 + \epsilon, 1)$ in the form $\phi = \sum_{t \in \{1,...,T\}} \epsilon^{t-1} \phi^t$ is to set T = 2, $\phi^1 = (1, 1)$, and $\phi^2 = (1, 0)$. The sequence (ϕ^1, ϕ^2) violates the eventual positivity requirement. By contrast, the weight vector $(1 + \epsilon, \epsilon) = (1, 0) + \epsilon(1, 1)$ is simple. The relevance of the distinction between simple and nonsimple hyperreal vectors, as well as the role played by the former, will become clear in Section 4.

Definition 2 (Simple hyperreal utilitarian welfare maximization (SHUWM)). A social welfare function W is a simple hyperreal utilitarian welfare function if

$$W(u) = \langle \phi, u \rangle = \sum_{t \in \{1, \dots, T\}} \epsilon^{t-1} \langle \phi^t, u \rangle,$$
(5)

where ϕ is a simple hyperreal weight vector. We say $u \in U$ maximizes a simple hyperreal utilitarian welfare function over U—or, more simply, u is a *SHUWM solution* of U—if there exists a simple hyperreal utilitarian welfare function W such that $W(u) \geq W(v)$ for all $v \in U$.

We can now state the first main result of the paper.

Theorem 1. Let U be a closed convex subset of \mathbb{R}^n and let u be a vector in U. Then, the following are equivalent:

- (i) u is Pareto optimal with respect to U.
- (ii) u is a SUWM solution of U.
- (iii) u is a SHUWM solution of U.

Proof. See Appendix A.

In the remainder of the section, we will offer interpretations of this result and insights into its proof.

We first start with the SUWM characterization of Pareto optimality implied by the equivalence between (i) and (ii). In SUWM, utilitarian welfare is maximized over multiple rounds for growing sets of agents until all agents are considered. From the social choice perspective, one can imagine a utilitarian social planner who prioritizes some agents—that is, those considered in earlier rounds of SUWM—and maximizes their (weighted) welfare before others. To achieve Pareto optimality, the social planner must assign *some* weights to all agents, but the welfare weights for some individuals (those who receive positive weights in later rounds) may need to be infinitely smaller than those for others (those who receive weights on individuals in different rounds). SUWM allows such flexibility by placing positive weights on individuals in different rounds. The eventual positivity condition encodes the requirement of Pareto optimality that "every agent's welfare counts" since the utility of each agent *i* has a positive weight in some round of welfare maximization.

The equivalence between (i) and (ii) is easy to visualize with the example in Figure 1, reproduced in Figure 4(a). In the first round, utilities are maximized within U with weights ϕ^1 , which is maximized by the thick vertical segment containing u. One can interpret this as the social planner first maximizing the utility of agent 1 while disregarding the welfare of the other individual completely. Since agent 1 is indifferent among all of these points, the social planner seeks to engage in further optimization. In the second (and last) round, utilities are again maximized but only within the vertical segment, now with an (arbitrary) nonnegative weight vector ϕ^2 that places a positive weight on agent 2. Hence, $\Phi = (\phi^1, \phi^2)$ is eventually positive. The weights ϕ^2 determine u as the unique maximizer, as illustrated in Figure 4(b). The theorem shows that the flexibility in assigning the weights in different rounds in SUWM enables an exact characterization.

The equivalence of (ii) and (iii) in the theorem shows that the sequential optimization involved in SUWM can be encoded in a one-shot weighted utilitarian welfare maximization with the introduction of simple hyperreal weights. This introduction of hyperreals allows for some agents to be prioritized over others in the sense of being assigned positive weights



Figure 4: Determining a Pareto optimal point in two rounds of sequential utilitarian welfare maximization.

in earlier rounds of SUWM. One can then interpret the former agents as carrying infinitely larger weights than the latter agents to constitute social welfare. The characterization in (iii) formalizes this idea by constructing the simple hyperreal weight vector $\phi = \sum_{t \in \{1,...,T\}} \epsilon^{t-1} \phi^t$. Since ϵ^s is infinitely larger than ϵ^t for any $t > s \ge 0$, the hyperreal vector ϕ assigns infinitely larger weights to the agents with higher priority than those with lower priority. For example, in Figure 4 the vector u = (1, 1) maximizes the simple hyperreal utilitarian welfare function with hyperreal weights $\phi^1 + \epsilon \phi^2 = (1 + \epsilon, \epsilon)$.⁹ While serving as a useful step toward our proof, this result lacks an important element that is fundamental in the economics context—that the weights be nonnegative and eventually positive. A nontrivial and crucial part of our proof lies in showing that nonnegative and eventually positive weights can be found if and only if the face consists of Pareto optimal points. The proof of Theorem 1 in Appendix A provides additional details and discussion.

Let us now explore some of the insights behind the proof of Theorem 1. The argument showing that (i) implies (ii) exploits a remarkable parallel between our problem and the question in convex geometry pertaining to *extreme faces* of a closed convex set. An extreme face, or simply a *face*, F of U is its convex subset whose elements cannot be expressed as convex combinations of points outside that set. (An extreme point is a special case of a face comprised of a singleton.) Geometrically, Pareto optimal points of U are made up of such faces (a result we establish). We say a hyperplane of U "exposes" a face F if it intersects U precisely at F, namely when F constitutes the set of points that maximize a linear function. A standard utilitarian welfare characterization of Pareto optima implies that the corresponding faces are "exposed" by hyperplanes with nonnegative weight vectors. From this perspective, the failure of standard weighted utilitarianism can be traced to the fact known in convex geometry that extreme faces may not always be exposed. However, an important finding in that literature is that an extreme face is "eventually exposed," that is, the face can be represented by the set of points that sequentially maximize *possibly*

⁹See Theorem 12.7 in Soltan (2015), reproduced as Lemma A.3 in the appendix.

negatively-weighted sum of utilities.¹⁰ While serving as a useful step toward our proof, this result lacks an element that is important for us and fundamental in the economics context— that the weights be nonnegative and eventually positive. A nontrivial and crucial part of our proof lies in showing that nonnegative and eventually positive weights can be found if and only if the face consists of Pareto optimal points.

Next, the fact that (ii) implies (iii) follows since any SUMW solution constitutes a SHUWM solution with the simple hyperreal weights constructed using a sequence of the SUWM weights as in (4). Finally, we establish that (iii) implies (i), by observing that any SHUWM solution must be Pareto optimal, given the positivity of the simple hyperreal weights.

4 Axiomatic foundation for "near" utilitarianism

In the previous section, we showed that "near" utilitarian welfare maximization—in the form of either SUWM or SHUWM—characterizes Pareto optima. Here we provide an axiomatic foundation for these welfare criteria. That is, we identify axioms of welfare orderings represented by these social welfare criteria.

This exercise serves at least two purposes. First, one can view the preceding characterization (Theorem 1) as providing a foundation for *some* version of utilitarianism. It is important to ask exactly what social welfare ordering corresponds to that version of utilitarianism. Second, our version of utilitarianism relaxes standard utilitarianism by allowing for a sequence of welfare weights or for hyperreal welfare weights in utilitarian welfare maximization. Identifying the social welfare orderings that justify such procedures will lay bare the precise nature of departure from those generating standard utilitarianism. This difference will in turn make precise, and flesh out, the sense in which our utilitarianism is "near" the standard one.

We begin with a state-of-the-art axiomatization of (weighted) utilitarianism. Let the social welfare ordering R^* be a complete and transitive binary relation defined over \mathbb{R}^n , the set of utility profiles of agents I, and let P^* and I^* denote the strict and indifferent parts of R^* , respectively. For any $u \in \mathbb{R}^n$ and any real number $\delta > 0$, let $B_{\delta}(u) := \{v \in \mathbb{R}^n : ||v - u|| < \delta\}$ be the δ -ball centered at u. Utilitarianism (with positive welfare weights) satisfies the following three axioms:

- Pareto Principle: for any u > v, we have uP^*v .
- Invariance: for any $u, v \in \mathbb{R}^n$, $a \in \mathbb{R}^n$ and $b \in \mathbb{R}_{++}$, if uR^*v , then $(a + bu)R^*(a + bv)$.
- Continuity: If uP^*v , then there exists $\delta > 0$ such that $u'P^*v$ for all $u' \in B_{\delta}(u)$.

Pareto Principle requires the welfare ordering to preserve the Pareto domination order. Invariance means that rescaling utility profiles by adding the same constant vector or by multiplying with the same positive coefficient does not alter their social welfare ordering. This

 $^{^{10}}$ See Theorem 12.7 in Soltan (2015), reproduced as Lemma A.3 in the appendix.

property permits just the right scope of interpersonal utility comparison that yields linear social welfare evaluation. Continuity means that perturbing the utilities of possibly all agents slightly does not alter social welfare ordering. Continuity forces welfare weights on alternative individuals to be of the same order of magnitude at the margin, meaning that no individual is treated infinitely better or worse compared with the others. Theorem 4.2-(2) of d'Aspremont and Gevers (2002) shows that utilitarianism is the only social welfare ordering that satisfies the three axioms:

Theorem 2. [D'Asprement-Gevers] Let R^* be a social welfare ordering. The following statements are equivalent:¹¹

- (i) R^* satisfies the Pareto Principle, Invariance, and Continuity,
- (ii) There exists $\phi \in \mathbb{R}^n_{++}$ such that uR^*v if and only if $\sum_{i \in I} \phi_i u_i \ge \sum_{i \in I} \phi_i v_i$.

It is easy to see that Continuity fails in our simple hyperreal utilitarian welfare function. Recall that in Figure 4, the Pareto optimum u = (1, 1) maximizes the simple hyperreal utilitarian welfare function $W(\cdot)$ with weights $(1 + \epsilon, \epsilon)$, where $\epsilon > 0$ is an infinitesimal. Hence, W(1, 1) > W(1, 1/2), for example. Yet, for any real number $\delta > 0$, $W(1 - \delta, 1 - \delta) < W(1, 1/2)$, so W fails Continuity. Indeed, it is well-known that lexicographic preference orderings cannot be represented by a continuous utility function (see, for instance, pages 46-7 of Mas-Colell, Whinston, and Green (1995)).

While continuity in its general form cannot be satisfied, the additional structure of our *near* utilitarianism may accommodate some weaker version of continuity. Indeed, we identify the precise form of weakening of Continuity compatible with our near utilitarianism. For each agent $i \in I$ and a real number $\delta > 0$, let $B^i_{\delta}(u) := \{v \in \mathbb{R}^n : |v_i - u_i| < \delta, v_j = u_j, \forall j \neq i\}$ be the δ -ball around u but only in the *i*-th coordinate. This notion allows us to define:

• Weak Continuity: for any uP^*v , there exist $i \in I$ and $\delta > 0$ such that $u'P^*v$ for all $u' \in B^i_{\delta}(u)$.

Weak Continuity requires the social welfare ordering to be robust to perturbations of only *some* individual agent's utility, and not necessarily to *all* possible perturbations of the utility profile, as required by Continuity. We next present the desired axiomatization of our "near" weighted utilitarian welfare functions. To this end, we adapt SUWM to welfare orderings in a natural way.

Definition 3. We say u sequentially utilitarian welfare dominates v according to Φ if u sequentially maximizes utilitarian welfare over $\{u, v\}$ according to Φ .^{12,13}

¹¹Theorem 4.2-(1) of d'Aspremont and Gevers (2002) gives the characterization with nonnegative welfare weights when Pareto is replaced with a weaker Pareto-like condition.

¹²In words, u sequentially utilitarian welfare dominates v, if there exists a sequence of eventually positive weight vectors $\Phi = (\phi^1, \phi^2, \dots, \phi^T)$ satisfying: either $\phi^t u = \phi^t v$ for all t or there exists $\tau \ge 1$ such that $\langle \phi^t, u \rangle = \langle \phi^t, v \rangle$ for all $t < \tau$ and $\langle \phi^\tau, u \rangle > \langle \phi^\tau, v \rangle$.

¹³Note that this ranking leads to a *rational* order, i.e., a binary relation that is reflexive, complete, and transitive. To see the transitivity (since the other properties are obvious), consider profiles u, v, and w such

Theorem 3. Let R^* be a social welfare ordering. The following statements are equivalent.

- (i) R^* satisfies the Pareto Principle. Invariance, and Weak Continuity,
- (ii) There exists a nonnegative and eventually positive sequence of weight vectors $\Phi = (\phi^1, \phi^2, ..., \phi^T)$ such that for any $u, v \in \mathbb{R}^n$, uR^*v if and only if u sequentially utilitarian welfare dominates v according to Φ .
- (iii) There exists a simple hyperreal weight vector $\psi \in ({}^*\mathbb{R}_{++})^n$ such that for any $u, v \in \mathbb{R}^n, uR^*v$ if and only if $\sum_{i \in I} \psi_i u_i \ge \sum_{i \in I} \psi_i v_i$.¹⁴

Proof. See Appendix B.

For (iii), the restriction to simple hyperreal utilitarian welfare functions is crucial. Recall that simplicity captures the eventual positivity of the weight vectors required in our SUWM, and this feature is essential for a hyperreal utilitarian welfare function to retain the weak continuity property. To see this, recall the non-simple weight vector $\psi := (1 + \epsilon, 1)$ with an infinitesimal $\epsilon > 0$ discussed in Section 3. The welfare function associated with this weight vector fails Weak Continuity. To see this, consider utility profiles u := (1,0) and v := (0,1). We have uP^*v because $\langle \psi, u \rangle = 1 + \epsilon > 1 = \langle \psi, v \rangle$. However, for any $i \in I$, real number $\delta > 0$, and $u' \in B^i_{\delta}(u)$ with u' < u, we have $\langle \psi, u' \rangle < 1 = \langle \psi, v \rangle$, so $u'P^*v$ does not hold, a violation of Weak Continuity. The reason for this difference is that this non-simple hyperreal function cannot be supported by a nonnegative and eventually-positive sequence of weight vectors required by SUWM.

By contrast, consider the simple hyperreal utilitarian welfare function $W(\cdot)$ with weights $(1+\epsilon,\epsilon)$ that exposes u in Figure 1. While W fails to be continuous, it is weakly continuous. Although W fails Continuity, it satisfies Weak Continuity. Recall W(u) > W(v), for u = (1, 1) and v = (1, 1/2). And, W(u') > W(v) for any $u' \in B^2_{\delta}(u)$ if $\delta \in (0, 1/2)$.

To visualize some of this discussion, Figure 5 illustrates the relationship between the axiomatizations of different notions of utilitarianism described in Theorems 2 and 3 and Proposition 2 (the last result is discussed in the next section).

5 Other characterizations of Pareto optimality

In this section, we discuss other characterizations of Pareto optima. As will be seen, these characterizations are not only related to our "near"-utilitarian welfare maximizations but they also highlight certain aspects of them and thus help to interpret and understand them. At the same time, we will show that they differ in their axiomatic properties from our "near"-utilitarian welfare characterizations. Our discussion will therefore illustrate that the

that u and v sequentially utilitarian welfare dominate v and w, respectively: that is, $\langle \phi^{\tau}, u \rangle > \langle \phi^{\tau}, v \rangle$ for some τ and $\langle \phi^{t}, u \rangle = \langle \phi^{t}, v \rangle$ for all $t < \tau$ while $\langle \phi^{\tau'}, v \rangle > \langle \phi^{\tau'}, w \rangle$ for some τ' and $\langle \phi^{t}, v \rangle = \langle \phi^{t}, w \rangle$ for all $t < \tau$. Then, letting $\tau'' = \min\{\tau, \tau'\}$, we have $\langle \phi^{\tau''}, u \rangle > \langle \phi^{\tau''}, w \rangle$ and $\langle \phi^{t}, u \rangle = \langle \phi^{t}, w \rangle$ for all $t < \tau''$, implying u sequentially utilitarian welfare dominates w.

¹⁴As with the order based on sequential utilitarian welfare domination, an order based on this ranking is also rational (as hyperreal numbers follow the same ordering system as real numbers).



Figure 5: Illustrating the axiomatizations of different notions of utilitarianism. The universe is the set of all social welfare orderings.

axiomatic properties of our "near"-utilitarian characterizations are special and not shared by other possible characterizations of Pareto optima.

Weighted utilitarianism with general hyperreal weights. As we discussed, the restriction to *simple* hyperreal utilitarian welfare functions disciplines them to resemble utilitarianism. At the same time, hyperreal utilitarian welfare maximization, with no restriction, also characterizes Pareto optimality.

Proposition 1. Let U be a closed convex subset of \mathbb{R}^n . Then, $u \in U$ is Pareto optimal if and only if

$$u \in \arg\max_{u' \in U} \langle \psi, u' \rangle,$$

for some weight vector $\psi = (\psi_i)_{i \in I} \in ({}^*\mathbb{R}_{++})^n$.

Proof. See Appendix C.1 in the Supplementary Appendix.

This proposition highlights the ability to assign an infinitely larger weight to one agent relative to another as a crucial feature that enabled SHUWM to characterize Pareto optimality. Compared with simple hyperreal utilitarian welfare functions, however, the class of general hyperreal utilitarian welfare functions is too large to be declared near-utilitarian. As we already saw, the class includes non-simple hyperreal functions that do not satisfy Weak Continuity, let alone Continuity.

Theorem 1 tells us that such non-simple functions are not needed for characterizing Pareto optimality. To illustrate their superfluity, recall the non-simple hyperreal vector $\psi = (1 + \epsilon, 1)$, where ϵ is a positive infinitesimal number. We can see that any such vector can be replaced by a simple hyperreal vector (which does satisfy Weak Continuity), in this particular case, a real vector, with no loss on the ability to characterize Pareto optima.¹⁵ We showed that the welfare function associated with $\psi = (1 + \epsilon, 1)$ fails Weak Continuity. Indeed, the next proposition shows that the class of hyperreal utilitarian welfare functions in Proposition 1 entails no restriction on social welfare orderings beyond the Pareto Principle and Invariance.

Proposition 2. Let R^* be a social welfare ordering. The following statements are equivalent.

- (i) R^* satisfies the Pareto Principle and Invariance.
- (ii) There exists a hyperreal weight vector $\psi \in ({}^*\mathbb{R}_{++})^n$ such that uR^*v if and only if $\sum_{i \in I} \psi_i u_i \ge \sum_{i \in I} \psi_i v_i$.

Proof. See Appendix C.2 in the Supplementary Appendix.

Figure 5 illustrates the differences in how general hyperreal utilitarian and other utilitarian welfare functions are axiomatized.

Sequential Nash bargaining. The second characterization is motivated by an institutional/behavioral implementation of Pareto optima. As is well known from the second fundamental welfare theorem, a Pareto optimal allocation, say in an exchange economy, can be implemented by a competitive equilibrium under a suitable endowment.¹⁶ In the same spirit, one may ask what institution implements a given Pareto optimum in a more general environment. Our SUWM characterization of Pareto optima allows one to envision sequential negotiations as fulfilling this goal. That is, any Pareto optimal outcome can be seen as emerging from a sequence of negotiations among individuals whose relative bargaining powers in round t are determined by the welfare weights ϕ^t in the corresponding round of SUWM characterization.

To be specific, suppose each agent has a disagreement utility, normalized as zero, that is less than any Pareto optimal utility—i.e., $u \gg 0$ for every $u \in U^P$. Consider a collection of bargaining units $\mathcal{I} = \{I^1, \ldots, I^T\}$ satisfying $I^{t-1} \subsetneq I^t$ for each $t = 2, \ldots, T$ and $I^T = I$. Imagine that the agents engage in a sequence of bargaining: in round 1, agents in I^1 bargain from U to a set $V^1 \subset U$, and in round $t = 2, \ldots, T$, agents in set I^t bargain from V^{t-1} to a set V^t . The bargaining protocol in each round t is a generalized Nash bargaining game (Kalai, 1977) in which each agent $i \in I^t$ has a bargaining power $\psi_i^t > 0$ such that $\sum_{i \in I^t} \psi_i^t = 1$ and a disagreement payoff 0. More specifically, for bargaining units $\mathcal{I} =$

¹⁵In this case, the real weight vector (1, 1) can be used in place of ψ in the sense that every Pareto optimal point that maximizes the hyperreal weight vector ψ also maximizes the real weight vector (1, 1).

 $^{^{16}}$ As an aside, in Appendix E in the SupplementaryAppendix, we illustrate how to use some of the techniques established in our proof of Theorem 1 to offer a new proof of the second welfare theorem that allows for weaker assumptions than the standard treatment. We discuss this more in the paper's conclusion section.

 $\{I^1, \ldots, I^T\}$ and bargaining powers $\Psi = (\psi^1, \ldots, \psi^T)$ satisfying the above requirement, we let $V^t := \arg \max_{u \in V^{t-1}} \prod_{i \in I^t} u_i^{\psi_i^t}$ for each $t = 1, \ldots, T$ with $V^0 := U$. Then, we call any $u \in V^T$ a sequential Nash bargaining solution (SNBS) over U for \mathcal{I} and Ψ , and call u an SNBS over U if there exist such \mathcal{I} and Ψ .

Observe now that SNBS implements the SUWM procedure for the logarithmic transforms of utilities. Namely, u is an SNBS over $U \subset \mathbb{R}^n_{++}$ if and only if $v := (\ln u_1, ..., \ln u_n)$ is an SUWM solution of $V := \{(\ln u'_1, ..., \ln u'_n) : (u'_1, ..., u'_n) \in U\}$. This connection also makes it clear that SNBS provides another characterization of Pareto optima.

Proposition 3. A vector $u \in U \cap \mathbb{R}^n_{++}$ is Pareto optimal if and only if u is an SNBS over U.

Proof. See Appendix C.3 in the Supplementary Appendix.

This result provides a behavioral interpretation of our near-weighted utilitarian welfare maximization. Despite this close connection, we will see that the SNBS characterization differs in the social welfare ordering it induces from our near-weighted utilitarian characterizations. To see this, we first define the welfare ordering induced by SNBS. We say u sequentially Nash welfare dominates v according to bargaining units \mathcal{I} and bargaining powers Ψ if u is an SNBS over $\{u, v\}$ for \mathcal{I} and Ψ .

Since SNBS implements the SUWM procedure for the logarithmic transforms of utilities, Theorem 3 implies that the following axiom would fulfill the same role as Invariance.

• Log Invariance: for any $u, v \in \mathbb{R}^n_{++}$, if uR^*v , then $u'R^*v'$ for any $u', v' \in \mathbb{R}^n_{++}$ such that, for some $a \in \mathbb{R}^n$ and $b \in \mathbb{R}_{++}$, $\ln u'_i = a_i + b \ln u_i$ and $\ln v'_i = a_i + b \ln v_i$ for all $i \in I$.

Combining this axiom with the Pareto Principle and Weak Continuity defined earlier, we obtain the following axiomatization of the welfare ordering based on SNBS.

Corollary 1. Let R^* be a social welfare ordering defined on \mathbb{R}^n_{++} . Then, the following statements are equivalent.

- (i) R^* satisfies the Pareto Principle, Log Invariance, and Weak Continuity.
- (ii) There exist bargaining units \mathcal{I} and bargaining powers Ψ such that for any $u, v \in \mathbb{R}^{n}_{++}$, $uR^{*}v$ if and only if u sequentially Nash welfare dominates v according to \mathcal{I} and Ψ .

Proof. See Appendix C.4 in the Supplementary Appendix.

In particular, this corollary implies that while SNBS characterizes Pareto optimality, the welfare orderings implied by the criterion depart further from utilitarianism than our "near"-utilitarian welfare criteria. While it shares the Pareto Principle and Weak Continuity, it generally fails Invariance.

Piecewise linear concave welfare function. Some readers may not like the sequentiality of SUWM or the use of hyperreal numbers in SHUWM. This observation leads to the question of whether it is possible to characterize Pareto optima by a one-shot maximization of a real-valued welfare function. For such a characterization, the welfare function cannot be weighted utilitarian. In particular, the function must be nonlinear. Can we achieve the characterization with minimal relaxation of the linearity? This motivates the following approach.

A social welfare function W is a piecewise linear concave (PLC) welfare function characterized by $(\psi^1, \psi^2, \ldots, \psi^t)$ if

$$W(v) = \min_{t \in \{1,\dots,T\}} \langle \psi^t, v \rangle, \tag{6}$$

where $\psi^t \in \mathbb{R}^n_+$ for each t. One candidate for the weight vectors $(\psi^1, \psi^2, \dots, \psi^T)$ to construct a PLC welfare function are those identified in the SUWM characterization; i.e., eventually positive weights. However, the characterization does *not* hold without an auxiliary condition. For this condition, let us say that a PLC welfare function W achieves its maximum over U via eventually positive weights if (i) $(\psi^1, \psi^2, \dots, \psi^T)$ is nonnegative and eventually positive and (ii) for all $v \in \arg \max_{u' \in U} W(u')$, $W(v) = \langle \psi^T, v \rangle$.

Proposition 4. Let U be a closed convex subset of \mathbb{R}^n . Then, $u \in U \cap \mathbb{R}^n_{++}$ is Pareto optimal if and only if it maximizes a PLC welfare function that achieves its maximum over U via eventually positive weights.^{17,18}

Proof. See Appendix C.5 in the Supplementary Appendix.

The role of the auxiliary condition is to prevent a Pareto suboptimal point from maximizing the PLC function (so that the "if" direction holds). To see it, observe that for any Pareto suboptimal point u, one can find v > u so that $W(v) \ge W(u)$. If u were a maximizer of W, then the auxiliary condition would require $W(v) = \langle \psi^T, v \rangle = \langle \psi^T, u \rangle = W(u)$ or $\langle \psi^T, v - u \rangle = 0$, which cannot hold since $\psi^T \gg 0$ and v > u. While achieving the goal of characterizing Pareto optima, the auxiliary condition also captures the main feature of SUWM that every agent's welfare must count as it requires a PLC function to be maximized via a weight vector that puts a positive weight on every agent's utility.

While our PLC welfare functions successfully characterize Pareto optima, we regard them to be further away from utilitarianism than our "near"-utilitarian welfare criteria. This is because, to our knowledge, no natural axioms characterize PLC welfare functions. In fact, it

¹⁷We focus on points $u \in \mathbb{R}^{n}_{++}$ for technical simplicity. This is not a substantive restriction because the economic environment is arguably unchanged when a constant is added to all utility profiles.

¹⁸This proposition may be reminiscent of construction of a PLC utility function based on an individual's choice data (see Afriat (1967)). The PLC social welfare function reveals the planner's preferences for agents' utilities similarly to how Afriat's PLC utility function reveals an individual's preferences for alternative goods. Note, however, that there are clear differences. The multiple linear components of our PLC welfare function result from multiple welfare weights corresponding to the successive rounds of SUWM. By contrast, the linear components in Afriat's construction reflect different budget lines a consumer faces in different choice scenarios. Moreover, the role played by the auxiliary condition to ensure every agent's welfare counts has no analogue in Afriat's characterization.

is not even obvious how to formulate a PLC function as a social welfare ordering in the face of the auxiliary condition. For instance, define the binary relation R^* by uR^*v if $W(u) \ge W(v)$ and $W(u) = \langle \psi^T, u \rangle$. Note that the condition $W(u) = \langle \psi^T, u \rangle$ is an adaptation of the auxiliary condition to the context of social welfare ordering. Then, R^* is not necessarily a complete binary relation, as the following example shows.

Example 1. Let there be two agents 1 and 2, T = 2, $\psi^1 = (1,0)$, $\psi^2 = (1,1)$, u = (1,1) and v = (0,0). Then, we have W(u) = 1 > 0 = W(v) while $W(u) = 1 < 2 = \langle \psi^2, u \rangle$, so neither uR^*v nor vR^*u holds. Hence, R^* is not complete.

6 Conclusion

We have provided two characterizations of Pareto optimal solutions of a closed convex set that are "near" to weighted utilitarian maximization in an axiomatic sense. They arise from relaxing the **Continuity** axiom that defines weighted utilitarian to a **Weak Continuity** axiom. We have shown that other characterizations of Pareto optimality are more "distant" from weighted utilitarianism because they violate more of its defining axioms. These results constitute significant progress in clarifying the connection between Paretian and utilitarian notions that are foundational to welfare economics.

Although our paper directly worked with the space of utility profiles U, our results drive implications for problems stated in the choice space X. Indeed, examining the structure of what points in the choice set give rise to Pareto optima has been a major focus in the multiobjective optimization literature. An early contribution in that literature is Charnes and Cooper (1967), who showed an equivalence between the problem of finding Pareto optimal solutions (in the choice set X) and that of solving a constrained nonlinear programming problem. Following their contribution, techniques in nonlinear programming were utilized to characterize Pareto optima under various conditions (Ben-Israel, Ben-Tal, and Charnes, 1977; Van Rooyen, Zhou, and Zlobec, 1994; Glover, Jeyakumar, and Rubinov, 1999; Ben-Tal, 1980) all of which require some form of differentiability of the utility functions. We believe further investigation into our approach may have the potential to add to this literature in at least two aspects. First, our characterization does not assume any form of differentiability. Indeed, the subtlety of non-exposure of Pareto optimal faces can also arise when utility functions are not smooth, as is often the case. Our methods may suggest ways to handle Pareto optimality when differentiability fails. Second, our methods may suggest a bridge between existing results in the choice space and results in the utility possibility space, where notions of (sequential) welfare maximization are salient and allow for more natural economic interpretations. Indeed, none of the characterizations in the above references speak to notions of welfare maximization.

A second area of future work would be to examine how the notion of exposure can be used to enhance separating hyperplane arguments that may arise in other economic settings. For instance, the second welfare theorem relies on the existence of a strictly positive weight vector for a supporting hyperplane (which constitutes equilibrium prices). One can prove this with a weaker assumption than in the existing proof of the theorem by leveraging the idea of exposing a Pareto optimal point—which is a target Pareto efficient allocation—, as we show in the Supplementary Appendix (see Appendix E). We believe there is scope to explore other economic settings where separating hyperplane arguments are used and similarly relax the conditions needed to ensure strict positivity when Pareto optimality (in combination with notions of exposure) may be used to assure the existence of a separating hyperplane with a positive weight vector.

A third area of future work is to extend the characterization presented in this paper to the case of infinite-dimensional economies. This is not a straightforward extension. Our argument in the finite-dimensional case depends on a termination condition that counts dimension. In the infinite-dimensional case, this termination condition is not accessible to us. Generalization would likely require a set convergence argument (for instance, using Hausdorff or Kuratowksi set-based metrics) that avoids discussion of dimension.

A Appendix: Proof of Theorem 1

A.1 Proof of (ii) \Rightarrow (iii)

Given the sequence $\Phi = (\phi^1, \ldots, \phi^T)$ that is sequentially maximized by u, let us construct a simple hyperreal weighted welfare function $W(\cdot)$ as in (5). Letting U^0, U^1, \ldots, U^T be the notation used in Definition 1, observe first that

$$W(v) = \langle \phi, v \rangle = \langle \phi, u \rangle = W(u) \text{ for every } v \in U^T,$$
(7)

by construction of ϕ and U^T . Next, consider any $v \notin U^T$. Then, there exists $\tau \in \{1, \ldots, T\}$ such that $\langle \phi^t, v \rangle = \langle \phi^t, u \rangle$ for all $t < \tau$ and $\langle \phi^\tau, v \rangle < \langle \phi^\tau, u \rangle$. Therefore,

$$\langle \phi, u \rangle - \langle \phi, v \rangle = \sum_{t=1}^{T} \epsilon^{t-1} \langle \phi^{t}, u \rangle - \sum_{t=1}^{T} \epsilon^{t-1} \langle \phi^{t}, v \rangle$$

$$= \epsilon^{\tau-1} \left(\langle \phi^{\tau}, u \rangle - \langle \phi^{\tau}, v \rangle \right) + \sum_{t \in \{\tau+1, \dots, T\}} \epsilon^{t-1} \left(\langle \phi^{\tau}, u \rangle - \langle \phi^{\tau}, v \rangle \right)$$

$$= \epsilon^{\tau-1} \left[\left(\langle \phi^{\tau}, u \rangle - \langle \phi^{\tau}, v \rangle \right) + \sum_{t \in \{\tau+1, \dots, T\}} \epsilon^{t-\tau} \left(\langle \phi^{\tau}, u \rangle - \langle \phi^{\tau}, v \rangle \right) \right]$$

$$(8)$$

where all arithmetic operations are valid because $*\mathbb{R}$ is an ordered field (Theorem 3.6.1 of Goldblatt (2012)). Because ϵ is an infinitesimal number strictly larger than zero, and $\langle \phi^{\tau}, u \rangle - \langle \phi^{\tau}, v \rangle$ is a positive real number, the expression inside the square bracket in (8) is positive, so $\langle \phi, u \rangle - \langle \phi, v \rangle > 0$ (see pages 50 and 51 of Goldblatt (2012)). Thus, we have shown that

$$W(v) = \langle \phi, v \rangle < \langle \psi, u \rangle = W(u) \text{ for every } v \notin U^T.$$
(9)

By equations (7) and (9), we have established that $u \in \arg \max_{v \in U} W(v)$.

A.2 Proof of (iii) \Rightarrow (i)

Suppose for contradiction that u maximizes the simple hyperreal weighted welfare function W as in (5) but is not Pareto optimal. Then, there exists $v \in U$ such that v > u. Since the sequence (ϕ^1, \ldots, ϕ^T) is nonnegative and eventually positive, we have $\phi_i = \sum_{t \in \{1, \ldots, T\}} \epsilon^t \phi_i^t > 0$ for each $i \in I$. Thus, $W(v) - W(u) = \langle \phi, v \rangle - \langle \phi, u \rangle = \sum_{i \in I} \phi_i(v_i - u_i) > 0$, a contradiction.

A.3 Proof of (i) \Rightarrow (ii)

We begin with some preliminaries before providing the proof in Appendix A.3.2.

A.3.1 Preliminaries

Let us first introduce a few concepts that are crucial for our analysis. A face of U is a nonempty convex subset F of U with the property that if $u \in F$ and $u = \alpha v + (1 - \alpha)w$ for some $0 < \alpha < 1$ and $v, w \in U$ then it must be that $v, w \in F$. That is, F is a face of a convex set if none of its elements are convex combinations of elements that lie outside of F. A proper face of U is a face of U that is a proper subset of U. A face F is an exposed face of U if there is a weight vector $\phi \in \mathbb{R}^n$ such that $F = \arg \max_{u \in U} \langle \phi, u \rangle$. In this case, we say that ϕ exposes F out of U. A face need not be exposed, as can be seen in Figure 1, where u is a singleton face that is not exposed.

For any convex subset G of U, its relative interior $\operatorname{ri}(G)$ is the set of all $u \in G$ such that for every $u' \in G$ there exists $\lambda > 0$ such that $u + \lambda(u - u') \in G$.

The following lemma shows a face structure of a convex set that is interesting in itself and useful for our analysis.

Lemma A.1 (Corollary 11.11(a) in Soltan (2015)). For a convex set $U \subseteq \mathbb{R}^n$, the collection of relative interiors of faces—that is, $\{ri(F) : F \text{ is a face of } U\}$ —forms a partition of U.

The next lemma shows that Pareto optimal points "come in faces." It is standard in the convex analytic literature to refer to Pareto optimal points as *maximal* points, so we use that language here.

Lemma A.2. Suppose a maximal point u of a closed convex set U lies in the relative interior of a face F of U. Then, every point in F is maximal.

Proof. The stated result is immediate in the case F is a singleton, so we may assume that F is not a singleton. Suppose for contradiction that F contains a nonmaximal element u'. Thus, there exists a $v \in U$ such that v > u'. Since $u \in ri(F)$, there exists $\lambda > 0$ such that $w' = u + \lambda(u - u') \in F$. Now let $z = \alpha w' + (1 - \alpha)v$, where $\alpha = \frac{1}{1+\lambda}$ or $\alpha(1 + \lambda) = 1$. Note that $z \in U$ since U is convex. Moreover,

$$z = \alpha \left(u + \lambda (u - u') \right) + (1 - \alpha)v = u - \alpha \lambda u' + (1 - \alpha)v = u + (1 - \alpha)(v - u') > u,$$

contradicting the maximality of u .

According to Lemma A.2, we say a face is *maximal* if all of its elements are maximal. Importantly for our purpose, Lemmas A.1 and A.2 imply that every maximal point of U belongs to a relative interior of a unique maximal face of U (possibly U itself).

The next result provides a key step of our argument: every face, possibly non-exposed, is *eventually* exposed.¹⁹

Lemma A.3 (Theorem 12.7 in Soltan (2015)). Let $U \subset \mathbb{R}^n$ be a convex set and F be a nonempty proper face of U. There is a sequence of convex sets $(G^t)_{t=0}^T$ such that

$$F = G^T \subset G^{T-1} \subset \dots \subset G^1 \subset G^0 = U,$$

where G^t is a nonempty proper exposed face of G^{t-1} for each $t = 1, \ldots, T$.

This lemma is already illustrated in the Introduction. In Figure 4, the singleton face u is exposed in two rounds: the vertical segment is exposed first by a weight vector (1, 0), and then u is exposed by weight vector (1, 1) (among many others) out of that vertical segment. This lemma is not enough for our result, however, as it is silent about any additional properties on the weight vectors that expose the sequence of faces. Crucially, our characterization requires the weight vectors to be nonnegative and eventually positive.

For these additional features, we need to introduce a set of analytical tools. Let J be any subset of the index set I and let χ^J denote the vector whose *i*-th coordinate is equal to 1 for every $i \in J$ and equal to 0 for every $i \notin J$. When J is the singleton $\{i\}$ we simplify $\chi^{\{i\}}$ to χ^i . A convex set U is downward closed in coordinates $J \subset I$ if, for all $u \in U$ and all $\tau \geq 0$, $u - \tau \chi^K \in U$ for any subset K of J. A convex set that is downward closed in all coordinates I is simply called downward closed. The downward closure of a closed convex set U is the downward closed set $dc(U) := \bigcup_{u \in U} (u - \mathbb{R}^n_+)$. It is straightforward to see that dc(U) is closed and convex if U is closed and convex.

One useful feature of downward closure is that it preserves maximal elements and thus maximal faces.

Lemma A.4. The set of maximal elements of a closed convex set coincides with that of its downward closure. If F is a maximal face of U then F is a maximal face of dc(U).

Proof. Let U be a closed convex set and dc(U) its downward closure. Let u be a maximal element of dc(U); that is, $(u + \mathbb{R}^n_+) \cap dc(U) = \{u\}$. If $u \in U$ then this implies $(u + \mathbb{R}^n_+) \cap U = \{u\}$ since $U \subset dc(U)$ and so u is a maximal element of U. Note that if $u \in dc(U) \setminus U$ then it cannot be maximal. Indeed, this implies that u = v - w for some $v \in U$ and nonzero $w \in \mathbb{R}^n_+$ and so v > u and so u is not maximal.

Conversely, we prove the contrapositive. Suppose $u \in dc(U)$ is not a maximal element. This implies that there exists a $w \neq u$ with $w \in dc(U)$ and $w \geq u$. However, then we can find a $v \geq w \geq u$ and $v \neq u$ and $v \in U$. This implies that u is not a maximal element of U. We next prove the second statement. To see that F is a face of dc(U), consider any $x, y \in dc(U)$ and $\lambda \in (0, 1)$ such that $z = \lambda x + (1 - \lambda)y \in F$. We need to show that both x and y belong

¹⁹Theorem 5 of Lopomo, Rigotti, and Shannon (2022) proves the same result for singleton faces F, i.e., extreme points.

to F. We first show that x and y are both maximal. Suppose for contradiction that x is not maximal. Then, we must have some $x' \in dc(U)$ such that x' > x. Let $z' = \lambda x' + (1 - \lambda)y$ and observe that $z' \in dc(U)$, $z' \ge z$, and $z' \ne z$, which contradicts the maximality of z. Given that x and y are both maximal, we must have $x, y \in U$ since there is no maximal point in $dc(U) \setminus U$. That F is a face of U then implies $x, y \in F$ as desired.

Crucially for our arguments, halfspaces of the form $\{u : \langle \phi, u \rangle \leq W_{\phi}\}$ that contain downward-closed sets must have nonnegative weight vectors.

Lemma A.5. Let U be a set that is downward closed in coordinates $J \subset I$. If U is contained in the halfspace $\{u \in \mathbb{R}^n : \langle \phi, u \rangle \leq W_{\phi}\}$, then $\phi_j \geq 0, \forall j \in J$.

Proof. Suppose for contradiction that $\phi_j < 0$ for some $j \in J$. Let v be an arbitrary element of U. Since U is downward closed in coordinates J, we also have $v - \lambda \chi^j \in U$ for any $\lambda \geq 0$, where χ^j is the unit vector with 1 in component j. However, observe that $\langle \phi, v - \lambda \chi^j \rangle =$ $\langle \phi, v \rangle - \lambda \langle \phi, \chi^j \rangle = \langle \phi, v \rangle - \lambda \phi_j$. But $\langle \phi, v \rangle - \lambda \phi_j \to \infty$ as $\lambda \to \infty$ since $\phi_j < 0$. This contradicts the fact that is contained $\{u \in \mathbb{R}^n : \langle \phi, u \rangle \leq W_\phi\}$.

Lemma A.6. Let F be a face of a closed convex set U that is downward closed in coordinates $J \subset I$. If ϕ exposes F out of U, then F is downward closed in coordinates $J \setminus \text{supp } \phi$.

Proof. Take any $j \in K := J \setminus \operatorname{supp} \phi$ and set $u' = u - \epsilon \chi^j$ for some $u \in F$ and $\epsilon > 0$. Since U is downward closed in coordinates J and $j \in J$, we have $u' \in U$. Moreover, $\langle \phi, u' \rangle = \langle \phi, u - \epsilon \chi^j \rangle = \langle \phi, u \rangle - \epsilon \langle \phi, \chi^j \rangle = \langle \phi, u \rangle - \epsilon \phi_j = \langle \phi, u \rangle$ since $\phi_j = 0$ when $j \in K$ since no element of K lies in supp ϕ . However, then $u' \in F$ since $\langle \phi, u' \rangle = \langle \phi, u \rangle = \max_{v \in U} \langle \phi, v \rangle$ and $F = \arg \max_{v \in U} \langle \phi, v \rangle$ since F is exposed by ϕ .

A.3.2 Proof of (i) \Rightarrow (ii)

Fix any maximal point u of U. We wish to show that u sequentially maximizes utilitarian welfare over U. The proof consists of several steps.

Step 1. There exists a unique face F of dc(U) such that $u \in ri(F)$. All points of F are maximal in dc(U).

Proof. By Lemma A.4, u is a maximal point of dc(U). By Lemma A.1 there is a unique face F of dc(U) which contains u in ri(F). By Lemma A.2, every point of F is maximal in dc(U), as desired.

Step 2. The face F (containing u) is a proper face of dc(U).

Proof. If not, we must have F = dc(U). Pick any $u' \in dc(U)$. Then, for any $\epsilon > 0$, $u'' = u' - \epsilon \chi^I$ is also in dc(U) by the downward closure property. Clearly, u'' is not a maximal point of dc(U) and cannot belong to F by Step 1, a contradiction.

Step 3. There exists a sequence of convex sets $(G^t)_{t=0}^T$ of dc(U) such that G^t is a proper exposed face of G^{t-1} for $t = 1, \ldots, T$, where $G^0 = dc(U)$, $G^T = F$, and $T \leq n$.

Proof. Since F is a proper face of dc(U) by Step 2, the result follows from Lemma A.3. For any set V, let $\dim(V)$ denote its dimension.²⁰ If V' is a proper face of convex set V, then $\dim(V') < \dim(V)$ by Theorem 11.4 in Soltan (2015). Thus, we have $T \leq n$ since $\dim(G^t) < \dim(G^{t-1})$ and since $\dim(G^0) = \dim(\operatorname{dc}(U)) = n$. \square

Step 4. There exists a sequence $\Phi = (\phi^1, \dots, \phi^T)$ such that for each $t = 1, \dots, T$, $G^t = \arg \max_{x \in G^{t-1}} \langle \phi^t, x \rangle,$

where $\phi^t > 0$ and $\operatorname{supp} \phi^1 \subset \operatorname{supp} \phi^2 \subset \ldots \subset \operatorname{supp} \phi^T = I.^{21}$

Proof. By Step 3, there exists a sequence of weight vectors $\Psi = (\psi^1, \ldots, \psi^T)$ such that, for each $t = 1, \ldots, T, \psi^t$ exposes G^t out of G^{t-1} . We construct $\Phi = (\phi^1, \ldots, \phi^T)$ with the stated properties.

The construction is recursive. First, since $G^0 = \operatorname{dc}(U)$, by Lemma A.5, $\phi^1 := \psi^1$ is nonnegative. For an inductive hypothesis, suppose that there are ϕ^k , $k = 1, \ldots, t - 1$, with the stated properties and that for each $k = 1, \ldots, t-1, G^k$ is downward-closed in coordinates $J^k := \{i \in I \mid \phi_i^k = 0\} = I \setminus \operatorname{supp} \phi^k$. Note that $J^{t-1} \subset J^{t-2} \subset \cdots \subset J^0 := I$. We will now construct ϕ^t and show that G^t is downward-closed in coordinates $J^t = \{i \in I \mid \phi_i^t = 0\}$.

First, observe G^{t-1} is contained in $\{u : \langle \psi^t, u \rangle \leq \max_{u' \in G^{t-1}} \langle \psi^t, u' \rangle \}$ and G^{t-1} is downwardclosed in coordinates J^{t-1} . Hence, Lemma A.5 implies that $\psi_i^t \ge 0$ on coordinates $j \in J^{t-1}$. Consider next $i \in \operatorname{supp} \phi^{t-1} = I \setminus J^{t-1}$. For such *i*, it is indeed possible for ψ_i^t to be negative. But noting $\phi_i^{t-1} > 0$ for such *i*, we define

$$\phi^t = \lambda^t \phi^{t-1} + \psi^t,$$

where $\lambda^t > \max_{i \in \text{supp } \phi^{t-1}} |\psi_i^t| / \phi_i^{t-1}$ is a (sufficiently large) positive scalar. Given this construction, $\phi_i^t \ge 0$ for all $i \in I$ and $\phi_i^t > 0$ for all $i \in \operatorname{supp} \phi^{t-1}$; i.e., $\operatorname{supp} \phi^t \supset \operatorname{supp} \phi^{t-1}$.

Let us show that ϕ^t exposes G^t out of G^{t-1} . To this end, let $M^t := \max_{x \in G^{t-2}} \langle \phi^{t-1}, x \rangle$. For all $x \in G^{t-1}$, we have

$$\langle \phi^t, x \rangle = \lambda^t \langle \phi^{t-1}, x \rangle + \langle \psi^t, x \rangle = \lambda^t M^t + \langle \psi^t, x \rangle,$$

since $\langle \phi^{t-1}, x \rangle = M^t$ for all $x \in G^{t-1}$. Henceforth,

$$\arg \max_{x \in G^{t-1}} \langle \phi^t, x \rangle = \arg \max_{x \in G^{t-1}} \langle \psi^t, x \rangle = G^t.$$

Since G^{t-1} is downward-closed in coordinates J^{t-1} and ϕ^t exposes G^t out of G^{t-1} , Lemma A.6 implies that G^t is downward-closed in coordinates $J^{t-1} \setminus \operatorname{supp} \phi^t = (I \setminus \operatorname{supp} \phi^{t-1}) \setminus \operatorname{supp} \phi^t =$ $I \setminus \operatorname{supp} \phi^t = J^t$, where the penultimate equality holds since $\operatorname{supp} \phi^{t-1} \subset \operatorname{supp} \phi^t$.

It remains to show that for each $i \in I$, there exists $t \in \{1, \ldots, T\}$ such that $\phi_i^t > 0$. To show this, it suffices to show that $\phi^T \gg 0$. Supposing not, there must be some $i \in I$ such that $\phi_i^t = 0$ for all $t = 1, \ldots, T$, so $i \in J^t$ for all $t = 1, \ldots, T$. Then, Lemma A.6 implies that for all t = 1, ..., T, G^t is downward-closed in coordinate *i*, which contradicts the fact that $G^T = F$ is maximal.

²⁰The dimension $\dim(V)$ of a convex subset V of U, including one of U's faces, is defined by the dimension of its affine hull: aff $(V) := \{\sum_{j=1}^{k} \alpha_j v^j \mid k \in \mathbb{N}, v^j \in V, \alpha_j \in \mathbb{R}, \sum_{j=1}^{k} \alpha_j = 1\}.$ ²¹Note that Φ , with these properties, is eventually positive since $\phi^T \gg 0$.

We have so far shown that u sequentially maximizes welfare over dc(U). We now prove the main result: u sequentially maximizes welfare over U. To this end, the following last step suffices.

Step 5. u sequentially maximizes utilitarian welfare over U.

Proof. Recall a sequence of weight vectors Φ from Step 4. Let U^0, U^1, \ldots, U^T be convex subsets of U such that, for each $t = 1, \ldots, T$, U^t is the face of U^{t-1} exposed by weight vector ϕ^t ; i.e.,

$$U^t = \arg \max_{x \in U^{t-1}} \langle \phi^t, x \rangle,$$

where $U^0 := U$. It suffices to prove that $U^T = F$, as this will prove that u sequentially maximizes utilitarian welfare over U.

To this end, it suffices to prove that $F \subset U^t \subset G^t$ for each $t = 0, \ldots, T$. We proceed inductively for the proof. First, note that the claim is trivially true for t = 0 because $U^0 := U \subset dc(U) := G^0$ and $F \subset U = U^0$ by definition. Now, suppose that the claim holds for t. We show (i) $F \subset U^{t+1}$ and (ii) $U^{t+1} \subset G^{t+1}$ as follows.

For (i), fix any point v in F. Then, since $F \subset G^{t+1}$ and ϕ^{t+1} exposes G^{t+1} out of G^t , we have $\langle \phi^{t+1}, v \rangle \geq \langle \phi^{t+1}, w \rangle$ for every $w \in G^t$. Because $U^t \subset G^t$ by the inductive assumption,

$$\langle \phi^{t+1}, v \rangle \ge \langle \phi^{t+1}, w \rangle \tag{10}$$

for every $w \in U^t$. Moreover, $v \in U^t$ by the assumption that $F \subset U^t$. This fact, combined with (10), implies that ϕ^{t+1} is maximized by v over U^t and so $v \in U^{t+1}$, since ϕ^{t+1} exposes U^{t+1} out of U^t . This holds for every $v \in F$ and so $F \subset U^{t+1}$, implying (i) holds for t+1.

As for (ii), fix any point v in U^{t+1} . By (i), we know that

$$\langle \phi^{t+1}, v \rangle = \langle \phi^{t+1}, w \rangle \tag{11}$$

for any $w \in F$, since U^{t+1} is exposed by ϕ^{t+1} and F is a subset of U^{t+1} . Moreover, by the definition of G^{t+1} and the fact that $F \subset G^{t+1}$ by construction, we know that

$$\langle \phi^{t+1}, w \rangle \ge \langle \phi^{t+1}, z \rangle \tag{12}$$

for any $w \in F$ and $z \in G^t$. Combining (11) and (12) implies that $\langle \phi^{t+1}, v \rangle \geq \langle \phi^{t+1}, z \rangle$ for any $z \in G^t$. This, and the fact that $v \in G^t$ (which immediately follows from $v \in U^{t+1} \subset U^t \subset G^t$), means that $v \in G^{t+1}$. Since this holds for any $v \in U^{t+1}$, we can conclude that $U^{t+1} \subset G^{t+1}$, so (ii) holds for t + 1.

This completes the induction and establishes the result.

B Appendix: Proof of Theorem 3

B.1 Proof of (i) \Rightarrow (ii)

We first show that if R^* satisfies the Pareto Principle and Invariance, then there exists a sequence $\Phi = (\phi^1, \phi^2, \dots, \phi^T)$ of nonnegative and eventually positive weight vectors such that for any $u, v \in \mathbb{R}^n$, uP^*v if u sequentially utilitarian welfare dominates v, that is,

$$\langle \phi^t, u \rangle > \langle \phi^t, v \rangle$$
 for some t and $\langle \phi^s, u \rangle = \langle \phi^s, v \rangle$ for all $s < t.$ (13)

We will then show if R^* satisfies Weak Continuity in addition, then (13) implies uP^*v . Thus, uP^*v if and only if u sequentially utilitarian welfare dominates v.

To do so, define $S := \{s \in \mathbb{R}^n : 0R^*s\}$ and $Q := \{s + p : s \in S, p \in \mathbb{R}^n, \text{ and } p \ll 0\}$. We now observe that Q is convex. To this end, let $q, q' \in Q$ and q'' = tq + (1-t)q' for any $t \in (0,1)$. Then, q = s + p and q' = s' + p' for some $s, s' \in S$ and $p, p' \ll 0$, and q'' = ts + (1-t)s' + tp + (1-t)p'. Then, by Invariance, we have $0R^*ts$, $0R^*(1-t)s'$, and $(1-t)s'R^*ts + (1-t)s'$. Thus, by transitivity, we have $0R^*ts + (1-t)s'$, i.e., $ts + (1-t)s' \in S$. Since $tp + (1-t)p' \ll 0$, we have $q'' \in Q$.

Letting \bar{Q} denote the closure of Q, \bar{Q} is also convex. Note that $0 \in S \subset \bar{Q}$ and that by the Pareto Principle, 0 is a maximal point of both S and \bar{Q} . Also, there is a maximal face $F \subset S$ with $0 \in F$. Letting $G^0 := \bar{Q}$, the same proof as Step 4 in the proof of Theorem 1 can be used to show there exists a sequence $\Phi = (\phi^1, \ldots, \phi^T)$ such that for each $t = 1, \ldots, T$,

$$G^{t} = \arg \max_{x \in G^{t-1}} \langle \phi^{t}, x \rangle, \tag{14}$$

where $\phi^t > 0$, supp $\phi^t \supseteq$ supp ϕ^{t-1} , $\phi^T \gg 0$, and $G^T = F$.

Claim 1. (13) implies uP^*v

Proof. Suppose for contradiction that there are some u, v for which (13) holds but vR^*u . Let w := u - v. Then, by Invariance, $0R^*w$, so $w \in S \subset \overline{Q} = G^0$. By the hypothesis, we have $\langle \phi^s, w \rangle = 0, \forall s < t$. Since $0 \in F = G^T$, (14) implies $w \in G^s, \forall s < t$, which in turn implies $\langle \phi^t, w \rangle \leq \langle \phi^t, 0 \rangle = 0$, or $\langle \phi^t, u \rangle \leq \langle \phi^t, v \rangle$. We thus have a contradiction.

Claim 2. uP^*v implies (13)

Proof. Let us first prove that $\langle \phi^s, u \rangle = \langle \phi^s, v \rangle$, $\forall s$ implies uI^*v . Suppose for a contradiction that uP^*v . By Weak Continuity, there are i and $\delta > 0$ such that $u'P^*v$ for all $u' \in B^i_{\delta}(u)$. We can then find a round t in which $i \in \operatorname{supp} \phi^t \setminus \operatorname{supp} \phi^{t-1}$. Since $\langle \phi^s, \chi^i \rangle = 0$, $\forall s < t$ and $\langle \phi^t, \chi^i \rangle > 0$ for the unit vector χ^i whose i-th component is equal to 1, we have $\langle \phi^s, u - \delta' \chi^i \rangle = \langle \phi^s, u \rangle = \langle \phi^s, v \rangle$, $\forall s < t$ and $\langle \phi^t, u - \delta' \chi^i \rangle < \langle \phi^t, u \rangle = \langle \phi^t, v \rangle$ for any $\delta' > 0$, which implies by the former statement that $vP^*(u - \delta'\chi^i)$, contradicting that $u'P^*v$ for all $u' \in B^i_{\delta}(u)$.

Since $\langle \phi^s, u \rangle = \langle \phi^s, u \rangle$, $\forall s$ implies uI^*v , uP^*v implies that there must be some t such that $\langle \phi^s, u \rangle = \langle \phi^s, v \rangle$, $\forall s < t$ and $\langle \phi^t, u \rangle \neq \langle \phi^t, v \rangle$. Since $\langle \phi^t, u \rangle < \langle \phi^t, v \rangle$ would imply vP^*u by Claim 1, we must have $\langle \phi^t, u \rangle > \langle \phi^t, v \rangle$ as desired.

B.2 Proof of (ii) \Rightarrow (iii)

Consider the sequence $(\phi^1, \phi^2, \dots, \phi^T)$ in (ii). Then, uP^*v is equivalent to (13), which implies

$$\sum_{i \in I} \psi_i u_i - \sum_{i \in I} \psi_i v_i = \epsilon^{t-1} \Big[\langle \phi^t, u - v \rangle + \sum_{s>t} \epsilon^{s-t} \langle \phi^s, u - v \rangle \Big] > 0, \tag{15}$$

where the inequality holds since the first term in the square bracket is a positive real and the second term is infinitesimal. Conversely, if $\sum_{i \in I} \psi_i u_i > \sum_{i \in I} \psi_i v_i$, then there must be some s such that $\langle \phi^s, u \rangle > \langle \phi^s, v \rangle$. Letting t be the smallest such s, we must have $\langle \phi^r, u \rangle = \langle \phi^r, v \rangle, \forall r < t$: else if $\langle \phi^r, u \rangle < \langle \phi^r, v \rangle$ for some r < t, then one can use a similar argument to (15) to obtain $\sum_{i \in I} \psi_i v_i > \sum_{i \in I} \psi_i u_i$, a contradiction.

B.3 Proof of (iii) \Rightarrow (i)

That the welfare function in (5)—or a social welfare ordering it represents—satisfies the Pareto Principle and Invariance is straightforward to check. To check that it satisfies Weak Continuity, consider any uP^*v so that W(u) > W(v). As argued before, there must be some t such that $\langle \phi^t, u \rangle > \langle \phi^t, v \rangle$ and $\langle \phi^s, u \rangle = \langle \phi^s, v \rangle, \forall s < t$. Pick any $i \in \operatorname{supp} \phi^t \setminus \operatorname{supp} \phi^{t-1}$. For sufficiently small $\delta > 0$ and all $u' \in B^i_{\delta}(u)$, we have $\langle \phi^t, u' \rangle > \langle \phi^t, v \rangle$ while $\langle \phi^s, u' \rangle = \langle \phi^s, v \rangle, \forall s < t$. This implies as desired that for all $u' \in B^i_{\delta}(u)$,

$$W(u') - W(v) = \epsilon^{t-1} \Big[\langle \phi^t, u' - v \rangle + \sum_{s>t} \epsilon^{s-t} \langle \phi^s, u' - v \rangle \Big] > 0,$$

where the inequality holds for the same reason as (15) holds.

References

- AFRIAT, S. N. (1967): "The construction of utility functions from expenditure data," *International Economic Review*, 8(1), 67–77. 17
- ANDERSON, R. M. (1991): "Non-standard analysis with applications to economics," Handbook of Mathematical Economics, 4, 2145–2208. 7
- ARROW, K., E. BARANKIN, AND D. BLACKWELL (1953): "Admissible points of convex sets," in *Contributions to the Theory of Games*, ed. by Kuhn, and Tucker, pp. 87–91. Princeton University Press. 2, 3, 4, SA.8
- BEN-ISRAEL, A., A. BEN-TAL, AND A. CHARNES (1977): "Necessary and sufficient conditions for a Pareto optimum in convex programming," *Econometrica*, pp. 811–820. 18
- BEN-TAL, A. (1980): "Characterization of Pareto and lexicographic optimal solutions," in Multiple Criteria Decision Making Theory and Application, pp. 1–11. Springer. 18
- BEWLEY, T. F. (2009): General Equilibrium, Overlapping Generations Models, and Optimal Growth Theory. Harvard University Press. 2, 4, SA.7
- BITRAN, G. R., AND T. L. MAGNANTI (1979): "The structure of admissible points with respect to cone dominance," *Journal of Optimization Theory and Applications*, 29(4), 573–614. 3
- BLUME, L. (1986): "Lexicographic refinements of Nash equilibrium," mimeo. SA.1, SA.3
- BLUME, L., A. BRANDENBURGER, AND E. DEKEL (1991): "Lexicographic probabilities and choice under uncertainty," *Econometrica*, pp. 61–79. 7, SA.1
- BRONSHTEIN, E., AND L. IVANOV (1975): "The approximation of convex sets by polyhedra," *Siberian Mathematical Journal*, 16(5), 852–853. SA.10
- BROWN, D. J., AND A. ROBINSON (1975): "Nonstandard exchange economies," *Econometrica*, pp. 41–55. 7
- CHARNES, A., AND W. W. COOPER (1967): Management Models and Industrial Applications of Linear Programming. Wiley. 18

- CHE, Y.-K., J. KIM, AND F. KOJIMA (2019): "Weak monotone comparative statics," arXiv:1911.06442 [econ.TH]. 6
- DANIILIDIS, A. (2000): "Arrow-Barankin-Blackwell theorems and related results in cone duality: A survey," in *Optimization*, pp. 119–131. Springer. 3
- D'ASPREMONT, C., AND L. GEVERS (2002): "Social welfare functionals and interpersonal comparability," Handbook of Social Choice and Welfare, 1, 459–541. 5, 12
- DILMÉ, F. (2022): "Lexicographic Numbers and Stability in Extensive Form Games," . 7
- GLOVER, B. M., V. JEYAKUMAR, AND A. M. RUBINOV (1999): "Dual conditions characterizing optimality for convex multi-objective programs," *Mathematical Programming*, 84(1), 201. 18
- GOLDBLATT, R. (2012): Lectures on the Hyperreals: An Introduction to Nonstandard Analysis. Springer. 7, 19
- HARSANYI, J. C. (1955): "Cardinal welfare, individualistic ethics, and interpersonal comparisons of utility," *Journal of Political Economy*, 63(4), 309–321. 2
- JOSHI, K. D. (1983): Introduction to General Topology. New Age International. SA.3
- KALAI, E. (1977): "Nonsymmetric Nash solutions and replications of 2-person bargaining," International Journal of Game Theory, 6, 129–133. 15
- LOPOMO, G., L. RIGOTTI, AND C. SHANNON (2022): "Detectability, duality, and surplus extraction," *Journal of Economic Theory*, 204, 1–37. 21
- MAS-COLELL, A., M. D. WHINSTON, AND J. R. GREEN (1995): *Microeconomic Theory*. Oxford University Press. 12
- MILGROM, P., AND C. SHANNON (1994): "Monotone comparative statics," *Econometrica*, 62, 157–180. 6
- SOLTAN, V. (2015): Lectures on Convex Sets. World Scientific. 10, 11, 20, 21, 23, SA.8, SA.9
- TOPKIS, D. M. (1998): Supermodularity and Complementarity. Princeton University Press. 6
- VAN ROOYEN, M., X. ZHOU, AND S. ZLOBEC (1994): "A saddle-point characterization of Pareto optima," *Mathematical Programming*, 67(1-3), 77–88. 18

Supplementary Appendices for: "Near" weighted utilitarian characterizations of Pareto optima

C Proofs for alternate characterization of Pareto optimality in Section 5

C.1 Proof of Proposition 1

The "only if" direction. Suppose $u \in U$ is Pareto optimal. Then, by Theorem 1, there is a sequence $\Phi = (\phi^1, \phi^2, \ldots, \phi^T)$ of $T \leq n$ nonnegative and eventually positive weight vectors such that u sequentially maximizes Φ . Defining $\psi := \sum_{t \in \{1,\ldots,T\}} \epsilon^t \phi^t$, we have $\psi_i > 0$ for each $i \in I$ since the vectors $\Phi = (\phi^1, \phi^2, \ldots, \phi^T)$ are nonnegative and eventually positive. Also, we have $u \in \arg \max_{v \in U} \langle \psi, v \rangle$ by Theorem 1.

The "if" direction. To show the contrapositive, assume that u is not Pareto optimal. Then there exists $v \in U$ such that v > u. Then, for any weight vector $\psi = (\psi_i)_{i \in I}$ with $\psi_i \in {}^*\mathbb{R}$ and $\psi_i > 0$ for each $i \in I$, $\langle \psi, v \rangle - \langle \psi, u \rangle = \sum_{i \in I} \psi_i (v_i - u_i) > 0$. This means that $u \notin \arg \max_{u' \in U} \langle \psi, u' \rangle$, as desired.

C.2 Proof of Proposition 2

The (ii) \Rightarrow (i) direction is obvious. To prove (i) \Rightarrow (ii), we adopt the proof approach of Theorem 1' of an unpublished work by Blume (1986) who studies an individual's decision under uncertainty.²² Suppose that the social welfare ordering R^* satisfies the Pareto Principle and Invariance.

Lemma C.1. Let $U' = \{u^1, u^2, \ldots, u^m\} \subset \mathbb{R}^n$ be a finite subset of utility profiles such that $uP^*\tilde{u}$ for some $u, \tilde{u} \in U'$. Then, there exists a nonnegative and non-zero weight vector $\phi^{U'} \in \mathbb{R}^n_+$ such that, for any $u, \tilde{u} \in U', uR^*\tilde{u}$ if and only if $\langle \phi^{U'}, u \rangle \geq \langle \phi^{U'}, \tilde{u} \rangle$.

Proof. We utilize the following fact:

Lemma C.2 (Lemma 7 of Blume (1986)). Let v^1, \ldots, v^K and w^{K+1}, \ldots, w^L be vectors in \mathbb{R}^n . Then, one of the following two statements holds.

1. There exists $x \in \mathbb{R}^n_+ \setminus \{0\}$ such that

 $\langle x, v^k \rangle > 0$ for all $k \in \{1, \dots, K\}$, and $\langle x, w^\ell \rangle = 0$ for all $\ell \in \{K+1, \dots, L\}$.

 $^{^{22}}$ Blume (1986) is superseded by the published version, Blume, Brandenburger, and Dekel (1991), although our proof is more closely related to the former.

2. There exist $y \in \mathbb{R}_+^K, z \in \mathbb{R}^{L-K}$ such that

$$\sum_{k=1}^{K} y_k v^k + \sum_{\ell=K+1}^{L} z_\ell w^\ell \le 0.$$

Moreover, if y = 0, then $\sum_{\ell=K+1}^{L} z_{\ell} w^{\ell} \neq 0$.

Let $v^k, k = 1, ..., K$ be the vectors of the form $v^k = u^k - \tilde{u}^k$ where $u^k, \tilde{u}^k \in U'$ and $u^k P^* \tilde{u}^k$, and $w^\ell, \ell = K + 1, ..., L$ be the vectors of the form $w^\ell = u^\ell - \tilde{u}^\ell$ where $u^\ell, \tilde{u}^\ell \in U'$ and $u^\ell I^* \tilde{u}^\ell$. It suffices to show that the case 1 of Lemma C.2 holds, as then the conclusion of Lemma C.1 holds when we set the solution x for the case 1 of Lemma C.2 as $\phi^{U'}$. To show this, we will show that the case 2 of Lemma C.2 does not hold. Suppose to the contrary that the case 2 of Lemma C.2 holds, with solution (y, z). We will obtain a contradiction.

Let $\tilde{z}_{\ell} := |z_{\ell}|$ and $\tilde{w}^{\ell} := sgn(z_{\ell})w^{\ell}$. Then $(y, \tilde{z}) > 0$ and $\sum_{k=1}^{K} y_k v^k + \sum_{\ell=K+1}^{L} \tilde{z}_{\ell} \tilde{w}^{\ell} \leq 0.^{23}$ Because v^k for each $k = 1, \ldots, K$ is of the form $u^k - \tilde{u}^k$ with $u^k P^* \tilde{u}^k$ while \tilde{w}^{ℓ} for each $\ell = K + 1, \ldots, L$ is of the form $u^{\ell} - \tilde{u}^{\ell}$ with $u^{\ell} I^* \tilde{u}^{\ell}$, it follows that

$$\sum_{k=1}^{K} y_k \tilde{u}^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell \tilde{u}^\ell \ge \sum_{k=1}^{K} y_k u^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell u^\ell.$$
(16)

Since R^* satisfies the Pareto Principle, this implies that

$$\left(\sum_{k=1}^{K} y_k \tilde{u}^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell \tilde{u}^\ell\right) R^* \left(\sum_{k=1}^{K} y_k u^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell u^\ell\right).$$
(17)

Meanwhile, since $u^k P^* \tilde{u}^k$ for each k and $u^{\ell} I^* \tilde{u}^{\ell}$ for each ℓ by assumption, by repeated applications of Invariance,²⁴ it follows that

$$\left(\sum_{k=1}^{K} y_k u^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell u^\ell\right) P^* \left(\sum_{k=1}^{K} y_k \tilde{u}^k + \sum_{\ell=K+1}^{L} \tilde{z}_\ell \tilde{u}^\ell\right),$$

if there exists k with $y_k > 0$, a contradiction to (17). If $y_k = 0$ for all k, then $\sum_{\ell=K+1}^{L} z_\ell w^\ell \neq 0$ by assumption, so we have by (16),

$$\sum_{\ell=K+1}^{L} \tilde{z}_{\ell} u^{\ell} < \sum_{\ell=K+1}^{L} \tilde{z}_{\ell} \tilde{u}^{\ell}$$

²³To see why $(y, \tilde{z}) > 0$, note that if $(y, \tilde{z}) = 0$, then y = 0 and $\sum_{\ell=K+1}^{L} z_{\ell} w^{\ell} = 0$, a contradiction to case 2 of Lemma C.2.

²⁴Specifically, for any $u, v, \tilde{u}, \tilde{v} \in \mathbb{R}^n$ with uR^*v and $\tilde{u}R^*\tilde{v}$ as well as $\alpha, \beta \in \mathbb{R}_+$, we have $(\alpha u + \beta \tilde{u})R^*(\alpha v + \beta \tilde{v})$, with R^* replaced with P^* if uP^*v and $\alpha > 0$. This is because $(\alpha u + \beta \tilde{u})R^*(\alpha v + \beta \tilde{u})$ and $(\alpha v + \beta \tilde{u})R^*(\alpha v + \beta \tilde{v})$ by Invariance, and hence by transitivity of R^* , the desired relationship follows (and the relation being strict if uP^*v and $\alpha > 0$).

holds.²⁵ Since R^* satisfies the Pareto Principle, this implies that

$$\left(\sum_{\ell=K+1}^{L} \tilde{z}_{\ell} \tilde{u}^{\ell}\right) P^* \left(\sum_{\ell=K+1}^{L} \tilde{z}_{\ell} u^{\ell}\right).$$
(18)

Meanwhile, recalling again $u^{\ell}I^*\tilde{u}^{\ell}$ for each $\ell = K + 1, ..., L$, and applying Invariance repeatedly, we have that

$$\left(\sum_{\ell=K+1}^{L} \tilde{z}_{\ell} u^{\ell}\right) I^* \left(\sum_{\ell=K+1}^{L} \tilde{z}_{\ell} \tilde{u}^{\ell}\right),$$

a contradiction to (18). This completes the proof.

Now we proceed to complete the theorem. To do so, we define

$$\mathcal{U} := \{ U' \subset \mathbb{R}^n : |U'| < \infty, \exists u, v \in U', uP^*v \},\$$

and, for each $u \in \mathbb{R}^n$, define the collection $U^u \subset \mathcal{U}$ by

$$\mathcal{U}^u := \{ U' \in \mathcal{U} : u \in U' \}.$$

Then, we consider a family V of collections defined by

$$V := \{ \mathcal{U}^u : u \in \mathbb{R}^n \}.$$

Now, let $u^1, u^2, \ldots, u^m \in \mathbb{R}^n$ and consider

$$\bigcap_{k=1}^m \mathcal{U}^{u^k}.$$

Note that $\bigcap_{k=1}^{m} \mathcal{U}^{u^k} \neq \emptyset$ since $\{u^1, u^2, \ldots, u^m\} \in U^{u^k}$ for all $k \in \{1, \ldots, m\}$, that is, the family V has the finite intersection property.

Now, we invoke the following fact:

Lemma C.3 (Proposition 3.6 of Joshi (1983)). A collection of sets has the finite intersection property if and only if there is a filter that is a superset of that collection.

The preceding argument and the claim imply that there exists a filter that is a superset of V. By Zorn's lemma, there exists an ultrafilter Ω that is a superset of the above filter, and hence a superset of V. This ultrafilter is clearly free, that is, the intersection of all sets in the collection Ω is empty: This is because all sets of the form \mathcal{U}^u is an element of Ω , and $\bigcap_{u \in \mathbb{R}^n} \mathcal{U}^u = \emptyset$.²⁶

Now, consider the set of functions from \mathcal{U} to \mathbb{R} . We say that two functions r and s are equivalent if $\{U' \in \mathcal{U} : r(U') = s(U')\}$ is in Ω . It is straightforward to show that this is an equivalence relation and the set \mathbb{R} of those equivalence classes is an ordered field which extends \mathbb{R} .²⁷ We call \mathbb{R} the set of hyperreal numbers. It is well known that addition and

²⁵To see this, suppose for contradiction that (16) holds as equality. Then it would imply $\sum_{\ell} \tilde{z}_{\ell} \tilde{w}^{\ell} = \sum_{\ell \neq \ell} z_{\ell} w^{\ell} = 0$, where the first equality follows from the definitions of \tilde{z}_{ℓ} and \tilde{w}^{ℓ} .

²⁶To show $\cap_{u \in \mathbb{R}^n} \mathcal{U}^u = \emptyset$, suppose for contradiction that $\cap_{u \in \mathbb{R}^n} \mathcal{U}^u$ is nonempty, so there exists $U' \in \bigcap_{u \in \mathbb{R}^n} \mathcal{U}^u$. Then, by definition U' is a finite subset of \mathbb{R}^n . So there exists $v \in \mathbb{R}^n$ such that $v \notin U'$. This implies $U' \notin U^v$, so $U' \notin \bigcap_{u \in \mathbb{R}^n} \mathcal{U}^u$, a contradiction.

 $^{^{27}}$ See Blume (1986) for proofs of this property as well as others in this paragraph.

multiplication defined on \mathbb{R} extend readily to \mathbb{R} by pointwise operations, while the orders \geq and > also extend in a similar manner. It is also standard to show that there exists an infinitesimal number in \mathbb{R} .

Now, let $\psi \in ({}^*\mathbb{R})^n$ be such that, for each $i \in I$, ψ_i is the equivalence class that contains the element r_i such that $r_i(U') = \phi_i^{U'}$ for each $U' \in \mathcal{U}$ and $\phi^{U'}$ given in Lemma C.1. Consider any $u, v \in \mathbb{R}^n$. We know that $\mathcal{U}^u \cap \mathcal{U}^v \in \Omega$. For any $U' \in \mathcal{U}^u \cap \mathcal{U}^v$, $u, v \in U'$, so if uR^*v , then $\langle \phi^{U'}, u \rangle \geq \langle \phi^{U'}, v \rangle$. By construction of ψ , this implies that $\langle \psi, u \rangle \geq \langle \psi, v \rangle$. A similar argument shows that uP^*v implies $\langle \psi, u \rangle > \langle \psi, v \rangle$. Finally, for each $i \in I$, note that $\chi^i P^*0$ as R^* satisfies the Pareto Principle. Therefore, it follows that $\psi_i = \langle \psi, \chi^i \rangle > \langle \psi, 0 \rangle = 0$, showing that $\psi \in ({}^*\mathbb{R}_{++})^n$. This completes the proof.

C.3 Proof of Proposition 3

Proof. For any $u \in \mathbb{R}^n_{++}$, let $\log u := (\log u_i)_i$ and, moreover, for any $u \in \mathbb{R}^n$, let $e^u := (e^{u_i})_i$. Let us also redefine $U := U \cap \mathbb{R}^n_{++}$ for notational simplicity. Now, let $\tilde{U} := \{\log u | u \in U\}$.

Claim 3. Suppose $u \in U$ and let $\tilde{u} = \log u$. Then, u is Pareto optimal with respect to U if and only if \tilde{u} is Pareto optimal with respect to $dc(\tilde{U})$.

Proof. First, note that $u \in U$ is Pareto optimal with respect to U if and only if \tilde{u} is Pareto optimal with respect to \tilde{U} because log(.) is a strictly increasing function. Second, note that $\tilde{u} \in \tilde{U}$ is Pareto optimal with respect to \tilde{U} if and only if it is Pareto optimal with respect to dc(\tilde{U}) because Pareto optimality is invariant to adding utility vectors to a set that are smaller than existing utility vectors. These two observations imply the conclusion of this claim.

Claim 4. Suppose that U is convex. Then $dc(\tilde{U})$ is convex.

Proof. Suppose $\tilde{u}, \tilde{u}' \in dc(\tilde{U})$, and $\lambda \in [0, 1]$. By definition of dc(.), it follows that there exist $\tilde{v}, \tilde{v}' \in \tilde{U}$ such that $\tilde{u} \leq \tilde{v}, \tilde{u}' \leq \tilde{v}'$. Therefore, by definition of \tilde{U} , there exist $v, v' \in U$ such that $\tilde{v} = \log v, \tilde{v}' = \log v'$.

Because U is convex, $w := \lambda v + (1 - \lambda)v'$ is in U. This implies that $\tilde{w} := \log w$ is in U. Now, because log(.) is a concave function, we have that

 $\lambda \tilde{v} + (1 - \lambda)\tilde{v}' = \lambda \log v + (1 - \lambda)\log v' \le \log(\lambda v + (1 - \lambda)v') = \log w = \tilde{w},$

so $\lambda \tilde{v} + (1-\lambda)\tilde{v}' \in \operatorname{dc}(\tilde{U})$. Because $\tilde{u} \leq \tilde{v}$ and $\tilde{u}' \leq \tilde{v}'$, it follows that $\lambda \tilde{u} + (1-\lambda)\tilde{u}' \in \operatorname{dc}(\tilde{U})$, as desired.

Now we proceed to prove the theorem.

The "if" direction: Suppose that $u \in U$ is an SNBS over U for some bargaining units \mathcal{I} and bargaining powers Ψ (satisfying the requirement). Then, $u \in V^T$ where $V^T = U$ and $V^t := \arg \max_{v \in V^{t-1}} \prod_{i \in I^t} v_i^{\psi_i^t}$ for each $t \geq 1$. This implies that $V^t := \arg \max_{v \in V^{t-1}} \sum_{i \in I^t} \psi_i^t \log v_i$. Setting $\tilde{u} := \log u$ and noting that ψ is a nonnegative and eventually positive sequence, \tilde{u} is a SUWM solution of dc(\tilde{U}) with respect to ψ . Therefore, by Theorem 1, \tilde{u} is Pareto optimal in dc(\tilde{U}). Then, by Claim 3, u is Pareto optimal with respect to U, as desired.

The "only if" direction: Suppose that $u \in U$ is Pareto optimal with respect to U. Then, by Claim 3, $\tilde{u} := \log u$ is Pareto optimal with respect to $dc(\tilde{U})$. Therefore, by Theorem 1, there exist a sequence $\phi := (\phi^t)_t$ of nonnegative and eventually positive welfare weight vectors such that $\tilde{u} \in \tilde{U}^T$ where $\tilde{U}^0 = dc(\tilde{U})$ and $\tilde{U}^t := \arg \max_{\tilde{u}' \in \tilde{U}^{t-1}} \sum_{i \in I^t} \phi_i^t \tilde{u}'_i$ for each $t \ge 1$. Then, for $u = e^{\tilde{u}}$, we have $u \in U^T$, where $U^t := \arg \max_{u' \in U^{t-1}} \prod_{i \in I^t} (u'_i)^{\phi_i^t} = \arg \max_{u' \in V^{t-1}} \prod_{i \in I^t} (u'_i)^{\psi_i^t}$ for each $t \ge 1$, where $V^t := \{e^{\tilde{v}} | \tilde{v} \in U^t\}$ and $\psi_i^t := \frac{\phi_i^t}{\sum_{j \in I^t} \phi_j^t}$, so u is an SNBS, as desired (note that ψ satisfies the condition required of bargaining powers for SNBS).

C.4 Proof of Corollary 1

To prove (ii) implies (i), we only check that R^* satisfies Log Invariance since the other axioms are rather straightforward to check. To do so, suppose that uR^*v so that for some $t \leq T$, $\prod_{i \in I^s} u_i^{\psi_i^s} = (>) \prod_{i \in I^s} v_i^{\psi_i^s}$ for s < (=)t + 1, which implies

$$\sum_{i \in I^s} \psi_i^s \ln u_i = (>) \sum_{i \in I^s} \psi_i^s \ln v_i \text{ for } s < (=)t + 1.$$
(19)

Consider now any u', v' such that for some $a \in \mathbb{R}^n$ and $b \in \mathbb{R}_{++}$, $\ln u'_i = a_i + b \ln u_i$ and $\ln v'_i = a_i + b \ln v_i$ for all $i \in I$. By (19), we have

$$\sum_{i \in I^s} \psi_i^s \ln u_i' = \sum_{i \in I^s} \psi_i^s a_i + b \left(\sum_{i \in I^s} \psi_i^s \ln u_i \right)$$
$$= (>) \sum_{i \in I^s} \psi_i^s a_i + b \left(\sum_{i \in I^s} \psi_i^s \ln v_i \right) = \sum_{i \in I^s} \psi_i^s \ln v_i' \text{ for } s < (=)t + 1,$$

which implies $\prod_{i \in I^s} (u'_i)^{\psi^s_i} = (>) \prod_{i \in I^s} (v'_i)^{\psi^s_i}$ for s < (=)t+1 or $u'R^*v'$ as desired.

We now prove that (i) implies (ii). Given any $u \in \mathbb{R}^n$, let e^u denote a vector $(e^{u_i})_{i \in I}$ and $\ln u$ denote a vector $(\ln u_i)_{i \in I}$ for simplicity. Consider any welfare ordering R^* on \mathbb{R}^n_{++} that satisfies the three axioms. Let us define another ordering \tilde{R}^* on \mathbb{R}^n as follows: for any $u, v \in \mathbb{R}^n$, $u\tilde{R}^*v$ if $e^uR^*e^v$. It is straightforward to check that \tilde{R}^* satisfies Pareto Principal, Invariace, and Weak Continuity. In particular, Invariance holds for the following reason. Consider any $u, v \in \mathbb{R}^n$ welfare or equivalently $\tilde{u} := e^uR^*e^v := \tilde{v}$. Invariance requires that for any $a \in \mathbb{R}^n$ and $b \in \mathbb{R}_{++}$, $u' := (a+bu)\tilde{R}^*(a+bv) := v'$ or equivalently $\tilde{u}' := e^{u'}R^*e^{v'} := \tilde{v}'$, which follows from $\tilde{u}R^*\tilde{v}$ and Log Invariance since $\ln \tilde{u}' = a + b \ln \tilde{u}$ and $\ln \tilde{v}' = a + b \ln \tilde{v}$. Since \tilde{R}^* satisfies the Pareto Principe, Invariance, and Weak Continuity, Theorem 3 implies that there exists a nonnegative and eventually positive sequence of weight vectors $\Phi = (\phi^1, \phi^2, ..., \phi^T)$ such that for any $u, v \in \mathbb{R}^n, u\tilde{R}^*v$ if and only if u sequentially utilitarian welfare dominates v according to Φ . For each t = 1, ..., T, let $I^t = \operatorname{supp} \psi^t$ and $\psi_i^t = \frac{\phi_i^t}{\sum_{i \in I^t} \phi_i^t}$ for all $i \in I^t$. Consider any u, v with uR^*v . Then, $\tilde{u} := \ln u\tilde{R}^* \ln v := \tilde{v}$ so that \tilde{u} sequentially utilitarian welfare dominates v according to $\Psi = (\psi^1, \ldots, \psi^T)$: that is, for some $t \leq T$, $\sum_{i \in I^s} \psi_i^s \tilde{u}_i = (>) \sum_{i \in I^s} \psi_i^s \tilde{v}_i$ for s < (=)t + 1, which implies that $\prod_{i \in I^s} u_i^{\psi_i^s} = (>) \prod_{i \in I^s} v_i^{\psi_i^s}$ for s < (=)t + 1, meaning u

sequentially Nash welfare dominates v according to \mathcal{I} and Ψ .

It is straightforward, and thus omitted, to prove that u sequentially Nash welfare dominating v according to \mathcal{I} and Ψ implies uR^*v .

C.5 Proof of Proposition 4

The "only if" direction: By Theorem 1, for any Pareto optimal $u \in U \cap \mathbb{R}_{++}^n$, there are nonnegative and eventually positive weights (ϕ^1, \ldots, ϕ^T) sequentially maximized by u. Letting U^t be defined as in (3), we have $u \in U^t$ for all $t = 1, \ldots, T$. Consider weights $(\psi^t)_{t=1}^T$ defined as $\psi^1 = \phi^1$ and $\psi^t = \frac{\langle \psi^{t-1}, u \rangle}{\langle \phi^t, u \rangle} \phi^t$ for each $t \geq 2$. First, using the fact that $u \in \mathbb{R}_{++}^n$ and $\phi^t \in \mathbb{R}_{+}^n, \forall t$, it is straightforward to see that $\langle \psi^t, u \rangle > 0$ and $\langle \phi^t, u \rangle > 0$ for every t. Thus, $\langle \psi^t, v \rangle \geq \langle \psi^t, v' \rangle$ if and only if $\langle \phi^t, v \rangle \geq \langle \phi^t, v' \rangle$ for all $v, v' \in U$. Note also that $\langle \psi^t, u \rangle = \frac{\langle \psi^{t-1}, u \rangle}{\langle \phi^t, u \rangle} \langle \phi^t, u \rangle = \langle \psi^{t-1}, u \rangle$ for each $t \geq 2$. Thus, we have $W(u) = \langle \psi^t, u \rangle$ for all $t = 1, \ldots, T$. Also, for any $v \in U^T$, we have $\langle \psi^t, u \rangle = \langle \psi^t, v \rangle$ for all t, so W(u) = W(v). For any $v \notin U^T$, there is some t such that $v \notin U^t$ so $\langle \psi^t, v \rangle < \langle \psi^t, u \rangle$, implying W(v) < W(u). Thus, u maximizes W, implying that W achieves its maximum over U via eventually positive weights.

The "if" direction: Consider any $u \in U$ maximizing a PLC function W that achieves its maximum via eventually positive weights. Suppose for contradiction that u is not Pareto optimal. Then, there is some v > u so that $\langle \psi^t, v \rangle \geq \langle \psi^t, u \rangle$ for all $t = 1, \ldots, T$. As u maximizes W, so does v. Given this and the fact that W achieves its maximum via eventually positive weights, we must have $\langle \psi^T, v \rangle = W(v) = W(u) = \langle \psi^T, u \rangle$ or $\langle \psi^T, v - u \rangle = 0$, which is a contradiction since $\psi^T \gg 0$ and v > u.

D Pareto optimality (U^P) and positive utilitarianism (U^{++})

This section aims to discover natural conditions for U^P to coincide with U^{++} . The following lemma, which follows easily from the proof of Theorem 1, is the key to our investigation.

Lemma D.1. If u is a maximal element of U that lies in the relative interior of an exposed face of dc(U) then u maximizes a positive weight vector over U.

Proof. In the proof of the "only if" part of Theorem 1 in Appendix A.3.2, if u is a maximal element of U that lies in the relative interior of an exposed face of dc(U), then T = 1 in Step 3 and by Step 4 we know ϕ^1 is positive. Hence, $\Phi = (\phi^1)$ and so by Step 5, we conclude that u maximizes the positive weight vector ϕ^1 over U.

To characterize when $U^P = U^{++}$, we need to introduce a few notions and establish their properties. First, the normal cone of U at a point $u \in U$ is the set

$$N_U(u) = \{ \phi \in \mathbb{R}^n \mid \langle \phi, u \rangle \ge \langle \phi, v \rangle \text{ for all } v \in U \}.$$

If $\phi \in N_U(u)$ then u is a maximizer of the linear function $\langle \phi, u \rangle$ over the set U. Then, the normal cone of a face $F \subset U$, denoted $N_U(F)$, as the normal cone of each of its relative interior points. Next, the relative boundary of F is defined as $F \setminus \operatorname{ri}(F)$.

The next two lemmas give us some properties of these notions.

Lemma D.2. Let F be a face of a convex set U. Then, every point in the relative interior of F has the same normal cone.

Proof. Let u, u' be distinct in the relative interior of F and suppose $N_U(u)$ contains an element ϕ not in $N_U(u')$. This implies $\langle \phi, u \rangle > \langle \phi, u' \rangle$. Since u is the relative interior, the point $v = u + \lambda(u - u')$ lies in F for a sufficiently small positive λ . However, $\langle \phi, v \rangle = \langle \phi, u \rangle + \lambda \langle \phi, u - u' \rangle > \langle \phi, u \rangle$, violating the assumption that ϕ is in $N_U(u)$. \Box

Lemma D.3. Let F be a face of a convex set U. Then, every point u in the relative boundary of F has $N_U(u) \supset N_U(F)$.

Proof. Let u be in the relative boundary of F. Suppose there is a weight vector ϕ in $N_U(v)$ (where v is any relative interior element of F) that is not in $N_U(u)$. That is,

$$\langle \phi, u \rangle \neq \langle \phi, v \rangle. \tag{20}$$

By the definition of the relative interior, we can get an element of the relative interior of F arbitrarily close to u, which yields a contradiction of the continuity of $\langle \phi, \cdot \rangle$ because of (20).

We are now ready to provide the condition that characterizes when $U^P = U^{++}$:

Proposition D.1. Let U be a closed convex set. Then $U^P = U^{++}$ if and only if every maximal element of U belongs to some exposed maximal face of dc(U).

Proof. The "if" direction. Observe that $U^{++} \subset U^P$ is immediate from Proposition 3.23 in Bewley (2009). It remains to show that $U^P \subset U^{++}$. Let $u \in U^P$. If u lies in the relative interior of an exposed face of dc(U), then $u \in U^{++}$ from Lemma D.1. The remaining case is where u lies on the relative boundary of a maximal exposed face F of dc(U). Since F is a maximal exposed face, then an element v in its relative interior maximizes a positive weight vector ϕ , again by Lemma D.1. By Lemma D.2, this implies that the normal cone $N_U(F)$ of face F contains ϕ and so, by Lemma D.3, the normal cone $N_U(u)$ of the point u contains ϕ . In other words, u maximizes the positive weight vector ϕ . This completes the proof.

The "only if" direction. Let u be a maximal element of U. By the equivalence of U^P and U^{++} , u maximizes a positive weight vector ϕ . Let $F = \arg \max_{v \in U} \langle \phi, v \rangle$. We claim that F is a maximal exposed face of dc(U), which contains u. The fact that F is maximal in dc(U) follows since Proposition 3.23 in Bewley (2009) (along with Lemma A.2) implies F is maximal in U and thus maximal in dc(U) by Lemma A.4. Suppose to the contrary that F is not exposed in dc(U). Then, there must exist an element $u' \in dc(U) \setminus U$ that maximizes ϕ but is not in F. However, since $u' \in dc(U) \setminus U$, there must exist $u'' \in U$ such that $u' \leq u''$ and $u'_i < u''_i$ for some index i. But this implies that $\langle \phi, u \rangle \geq \langle \phi, u'' \rangle > \langle \phi, u' \rangle$, where the weak inequality holds by the definition of F and the strict inequality holds since ϕ is positive. This yields a contradiction and so we conclude that F is an exposed face of dc(U).



Figure 6: The maximal extreme point u is not exposed while $U^P = U^{++}$.

We now discuss a few of the nuances in the statement of Proposition D.1. First, the condition cannot be weakened so that every maximal element of U simply lies in a (potentially nonmaximal) exposed face of dc(U). Consider our canonical example in Figure 1. The point u lies on an exposed face of dc(U), but this face is not a maximal face of dc(U).

Figure 1 also demonstrates that it is not sufficient for a point to lie on a maximal exposed face of U (as opposed to dc(U)) to guarantee it maximizes a positive weight vector. Consider the point u'', which is a maximal exposed extreme point of U but does not maximize any positive weight vector over U. However, u'' does not lie on a maximal exposed face of dc(U)and so does not contradict the theorem.

Given the above nuance, a simpler sufficient condition may be useful. Consider the setting where all maximal faces of dc(U) are exposed.

Corollary D.1. If U is a closed convex set such that all maximal faces of dc(U) are exposed, then $U^P = U^{++}$.²⁸

Proof. Note that every maximal element of U lies in a maximal face of dc(U) by Lemma A.4. This and the hypothesis imply that every maximal element of U belongs to some exposed maximal face of dc(U). Applying Proposition D.1, we obtain the desired conclusion.

However, the converse of Corollary D.1 is false, as illustrated by the example in Figure 6. One sufficient condition for the hypothesis of Corollary D.1 to hold is that U is a polyhedron. In that case, all faces of U are exposed (Theorem 13.21 of Soltan (2015)); moreover, its downward closure of a polyhedron is also a polyhedron (Theorem 13.20 of Soltan (2015)), so all of its faces are exposed.

Let X be a polyhedral subset of \mathbb{R}^m_+ (possibly \mathbb{R}^m_+ itself). The utility function $u_i : X \to \mathbb{R}$ is *piecewise-linear concave (PLC)* if there exist finite index set K_i and affine functions

²⁸This cannot be derived easily from Arrow, Barankin, and Blackwell (1953). To see this, recall that they establish $U^{++} \subset U^P \subset \operatorname{cl}(U^{++})$. This implies that if U^{++} is closed then $U^P = U^{++}$. However, in the "tilted cone" in Figure 2, U^{++} is not closed since the point K does not lie in U^{++} but is the limit point of elements in U^{++} (indicated by the line in the figure). However, it is straightforward to check that U^P and U^{++} coincide. One can also check that all maximal faces of dc(U) for U in Figure 2 are exposed, the condition of Corollary D.1.

 $u_{i,k}: \mathbb{R}^m_+ \to \mathbb{R}$ for each $k \in K_i$ such that $u_i(x) = \min_{k \in K_i} u_{i,k}(x)$ for all $x \in X$. The lemma uses some of the following facts.

Lemma D.4. The following properties on polyhedra hold:

- (i) Let P_1, P_2, \ldots, P_n be a finite collection of polyhedra in \mathbb{R}^m . The Cartesian product $P_1 \times P_2 \times \cdots \times P_n$ is a polyhedron in \mathbb{R}^{mn} .
- (ii) Let $\pi : \mathbb{R}^d \to \mathbb{R}^m$ be an affine map and let P be a polyhedron in \mathbb{R}^d . Then $\pi(P)$ is a polyhedron.
- (iii) All faces of a polyhedron are exposed.
- (iv) The downward closure of a polyhedron is also a polyhedron.

Proof. (i) Consider two polyhedra in \mathbb{R}^m , P_1 and P_2 . Letting $Q_1 := P_1 \times \mathbb{R}^m$ and $Q_2 := \mathbb{R}^m \times P_2$, each Q_k is a polyhedron in \mathbb{R}^{2m} , so $P_1 \times P_2 = \bigcap_{k=1,2} Q_k$ is a polyhedron in \mathbb{R}^{2m} . The result follows from applying this argument repeatedly. (ii) This is Theorem 13.21 in Soltan (2015). (iii) This is Corollary 13.12 in Soltan (2015). (iv) This follows by nothing that since $dc(P) = P + \mathbb{R}^n_-$ where \mathbb{R}^n_- is the nonpositive orthant of \mathbb{R}^n and by applying Theorem 13.20 of Soltan (2015).

Lemma D.5. If each agent has a PLC utility function defined on a polyhedron X and U is defined according to (1), then dc(U) is a polyhedron.

Proof. For each $k \in K_i$, let $X_{i,k} = \{x \in X \mid u_{i,k}(x) \leq u_{i,k'}(x), \forall k' \in K_i\}$. Since X is a polyhedron and all functions $(u_{i,k})_{k \in K_i}$ are affine, $X_{i,k}$ is an intersection of finitely many polyhedra and thus a polyhedron.

Now let $\mathcal{K} = \{\mathbf{k} = (k_i)_{i \in I} \mid k_i \in K_i \text{ for all } i\}$. For each $\mathbf{k} \in \mathcal{K}$, let $X_{\mathbf{k}} = \bigcap_{i \in I} X_{i,k_i}$ and observe that $X_{\mathbf{k}}$ is a polyhedron. Also, all functions $u_1(\cdot), \ldots, u_I(\cdot)$ are affine on $X_{\mathbf{k}}$ since for each $i \in I$, $u_i(x) = u_{i,k_i}(x), \forall x \in X_{\mathbf{k}}$. Then, by Lemma D.4(ii), the set $U_{\mathbf{k}} = \{(u_i(x))_{i \in I} \mid x \in X_{\mathbf{k}}\}$ is a polyhedron. Observe that $U = \{(u_i(x))_{i \in I} \mid x \in X\} = \bigcup_{\mathbf{k} \in \mathcal{K}} U_{\mathbf{k}}$. While we do not know whether the set U, which is a union of polyhedra, is a polyhedron, Theorem 13.19 of Soltan (2015) shows that $\overline{U} := cl(conv \cup_{\mathbf{k} \in \mathcal{K}} U_{\mathbf{k}})$ is a polyhedron, where cl and conv denote the closure and convex hull, respectively.

Next, we show that $\operatorname{dc}(U) = \operatorname{dc}(\overline{U})$. By definition of \overline{U} , $\operatorname{dc}(U) \subset \operatorname{dc}(\overline{U})$ is clear. To show $\operatorname{dc}(\overline{U}) \subset \operatorname{dc}(U)$, consider any $\tilde{u} \in \operatorname{conv} \cup_{\mathbf{k} \in \mathcal{K}} U_{\mathbf{k}}$ so that $\tilde{u} = \sum_{\mathbf{k} \in \mathcal{K}} \lambda_{\mathbf{k}} \tilde{u}_{\mathbf{k}}$ for some weight $(\lambda_{\mathbf{k}})_{\mathbf{k} \in \mathcal{K}}$ and $\tilde{u}_{\mathbf{k}} \in \bigcup_{\mathbf{k}' \in \mathcal{K}} U_{\mathbf{k}'}$. Also, for each $\tilde{u}_{\mathbf{k}}$, we can find $\tilde{x}_{\mathbf{k}} \in X_{\mathbf{k}}$ such that $(u_i(\tilde{x}_{\mathbf{k}}))_{i \in I} = \tilde{u}_{\mathbf{k}}$. Letting $x = \sum_{\mathbf{k} \in \mathcal{K}} \lambda_{\mathbf{k}} \tilde{x}_{\mathbf{k}}$, observe that $x \in X$ by the convexity of X and that for all $i \in I$, $u_i(x) \geq \sum_{\mathbf{k} \in \mathcal{K}} \lambda_{\mathbf{k}} u_i(\tilde{x}_{\mathbf{k}}) = \tilde{u}_i$ by the concavity of $u_i(\cdot)$, which means that $\tilde{u} \in \operatorname{dc}(U)$. Thus, $\operatorname{conv} \cup_{\mathbf{k} \in \mathcal{K}} U_{\mathbf{k}} \subset \operatorname{dc}(U)$, implying that $\operatorname{cl}(\operatorname{conv} \cup_{\mathbf{k} \in \mathcal{K}} U_{\mathbf{k}}) \subset \operatorname{dc}(U)$ since $\operatorname{dc}(U)$ is closed, from which $\operatorname{dc}(\overline{U}) \subset \operatorname{dc}(U)$ follows, as desired.

Lastly, observe that $dc(\overline{U}) = \overline{U} + \mathbb{R}^n_-$ and that both \overline{U} and \mathbb{R}^n_- are polyhedra, which implies (by Lemma D.4(iv)) that $dc(\overline{U}) = dc(U)$ is a polyhedron.

The following is obtained immediately from Corollary D.1 and Lemma D.5, and the fact that all faces of polyhedra are exposed. It is a clean economic setting where U^P and U^{++} coincide.

Proposition D.2. If each agent has a PLC utility function defined on a polyhedron X and U is defined according to (1), then $U^P = U^{++}$.²⁹

E Second welfare theorem with piecewise-linear concave utility functions

In the paper, we showed that the notions of exposed faces and normal vectors play crucial roles for our characterization of a Pareto optimal utility profile as a welfare-maximizing point. Recall that the normal vector also plays an important role in the second theorem of welfare economics in identifying a price vector that supports a Pareto optimal allocation as a competitive equilibrium outcome. Unlike in our characterization, the idea of a normal vector in the second welfare theorem applies to the space of goods, not the space of utility profiles. However, the fact that the two spaces are closely connected hints at the possibility of establishing the second welfare theorem using the machinery we have developed so far. We do so in the current section under a set of assumptions on the agent preferences and endowments that generalize the existing welfare theorem in a certain direction.

To begin, consider an exchange economy with m types of goods with some integer m > 0. For each $k \in \{1, \ldots, m\}$, let $\bar{e}^k > 0$ be the total supply of type-k goods in the environment. Let \bar{e} denote the vector $(\bar{e}^k)_{k=1}^m$. Each alternative $x = (x_i)_{i \in I}, x_i = (x_i^k)_{k=1}^m \in \mathbb{R}_+^m$, specifies consumption bundle x_i for each $i \in I$. A profile of consumption bundles x is said to be feasible if and only if $\sum_{i \in I} x_i \leq \bar{e}$. In this context, the choice set X is defined as the set of all feasible profiles of consumption bundles. Each individual $i \in I$ is endowed with a utility function $u_i : \mathbb{R}_+^m \to \mathbb{R}$. Suppose that each agent i is endowed with a vector of goods $e_i \in \mathbb{R}_+^m \setminus \{0\}$ and let $\bar{e} = \sum_{i \in I} e_i$. A vector $p \in \mathbb{R}^m$ is referred to as a price profile. A pair (p, x) of a price profile p and a profile $x = (x_i)_{i \in I}$ of consumption bundles is a Walrasian equilibrium if

- 1. $\sum_{i \in I} x_i = \bar{e}$, and
- 2. $x_i \in \arg \max_{y_i \in B_i(p)} u_i(y_i)$ for each $i \in I$, where $B_i(p) := \{y_i \in \mathbb{R}^m_+ \mid \langle p, y_i \rangle \leq \langle p, e_i \rangle\}$ is the budget set of i.

We consider a case where utility functions of all players are piecewise-linear concave (PLC), as defined in Appendix D. PLC utility functions may appear somewhat restrictive, but any concave function can be approximated arbitrarily closely by a PLC utility function (Bronshtein and Ivanov, 1975). Meanwhile, we make a weaker assumption in another dimension—preference monotonicity. The existing second welfare theorem assumes agents' utility functions to be strictly monotonic. We invoke a weaker form of monotonicity. Say that an allocation $(x_i)_{i\in I}$ is strictly feasible for good k if it is feasible and satisfies $\sum_{i\in I} x_i^k < \bar{e}^k$. We assume that the agent preferences are monotonic under limited resources in the following

²⁹It is worth noting that the ABB theorem provides an alternative proof of this result. Recall that it suffices to argue U^{++} is closed in order to conclude $U^P = U^{++}$. Clearly, the elements of U^{++} come in faces, and a polyhedron has finitely many faces. Since the faces of a polyhedron are closed, and a finite union of closed sets is closed, this implies that U^{++} is closed.

sense: for any allocation $(x_i)_{i \in I}$ that is strictly feasible for good k, there exist an agent j and $\tilde{x}_j \in \mathbb{R}^m_+$ such that $u_j(\tilde{x}_j) > u_j(x_j)$ while $\tilde{x}_j^{k'} = x_j^{k'}, \forall k' \neq k, \tilde{x}_j^k > x_j^k$, and $\tilde{x}_j^k + \sum_{i \neq j} x_j^k \leq \bar{e}^k$. That is, given any allocation that does not exhaust the endowment of good k, there exists an agent who gets better off by consuming more of that good within its endowment. This condition is fairly weak. For instance, it allows for agents to consider a certain good indifferently, or even as bads (rather than goods), as long as there is at least one agent who likes to consume that good. We are now ready to prove the second welfare theorem under the above assumptions.

Proposition E.1. Consider the exchange economy described above. If $(u_i(e_i))_{i \in I}$ is Pareto optimal, then there exists a positive price vector $p \gg 0$ such that $(p, (e_i)_{i \in I})$ is a Walrasian equilibrium.

Proof of Proposition E.1. Let $A_i := \{x \in \mathbb{R}^m_+ \mid u_i(x) \ge u_i(e_i)\}$ for each agent *i*. Observe that each A_i is a polyhedron since it is an intersection of two polyhedra, $\{x \in \mathbb{R}^m \mid x \ge 0\}$ and $\{x \in \mathbb{R}^m \mid u_i(x) \ge u_i(e_i)\} = \bigcap_{k \in K_i} \{x \in \mathbb{R}^m \mid u_{i,k}(x) \ge u_i(e_i)\}.$

Consider the set $A = \{x \in \mathbb{R}^m_+ \mid \exists x_1 \in A_1, x_2 \in A_2, \dots, x_n \in A_n \text{ s.t. } x = \sum_{i \in I} x_i\}$. Observe that A is the image of the set $A_1 \times A_2 \times \cdots \times A_n$ under the affine mapping π that maps $(x_i)_{i \in I}$ to $\sum_{i \in I} x_i$. By Lemma D.4(i) and (ii), A itself is a polyhedron.

Next, we argue that \bar{e} is a minimal element of the set A. Suppose for contradiction that there exists an element $x \in A$ where $x < \bar{e}$ where $x^k < \bar{e}^k$ for some good k. Since $x \in A$, there exists an allocation $(y_i)_{i\in I}$ where $y_i \in A_i$ such that $x = \sum_{i\in I} y_i$. Since this allocation is strictly feasible for the good k, the monotone preference under limited resources implies that there are some agent j and $\tilde{y}_j \in \mathbb{R}^m_+$ such that $u_j(y_j) < u_j(\tilde{y}_j)$ while $\tilde{y}_j^{k'} = y_j^{k'}, \forall k' \neq k$, $\tilde{y}_j^k > y_j^k$, and $\tilde{y}_j^k + \sum_{i\neq j} y_i^k \leq \bar{e}^k$. Now consider an alternative allocation $(z_i)_{i\in I}$, which is identical to $(y_i)_{i\in I}$ except that $z_j = \tilde{y}_j$. Note that this allocation is feasible under the endowment \bar{e} and that $u_j(z_j) > u_j(y_j) \geq u_j(e_j)$ while $u_i(z_i) = u_i(y_i) \geq u_i(e_i), \forall i \neq j$, which contradicts the Pareto optimality of $(e_i)_{i\in I}$.

That \bar{e} is a minimal element of A implies that $-\bar{e}$ is a maximal element of -A. By Lemma A.4, this implies that $-\bar{e}$ is a maximal element of dc(-A). Moreover, by Lemma D.4(iv) dc(-A) is a polyhedron and so by Lemma D.4(iii) all of its faces are exposed. Thus, by Lemma D.1, there exists a supporting hyperplane of -A through the point $-\bar{e}$ with a positive normal ϕ . The same normal $p := \phi$ can define a supporting hyperplane to A through the point \bar{e} ; that is,

$$\langle p, y \rangle \ge \langle p, \bar{e} \rangle, \forall y \in A,$$

where p is a positive vector of prices.

It remains to show that the positive price vector p just constructed supports the allocation $(e_i)_{i \in I}$ as a Walrasian equilibrium. For this, it suffices to show that each e_i maximizes $u_i(\cdot)$ under the prices p and the budget $\langle p, e_i \rangle$. To do so, we take any x_i with $u_i(x_i) > u_i(e_i)$ and show that agent i cannot afford x_i .

By continuity of u_i , the inequality $u_i(x_i) > u_i(e_i)$ implies that for some $\lambda < 1$ but sufficiently close to 1, we have $u_i(\lambda x_i) > u_i(e_i)$, so by definition we have $\lambda x_i \in A_i$. This implies that $\lambda x_i + \sum_{j \neq i} e_j \in A$. Since $\langle p, \lambda x_i + \sum_{j \neq i} e_j \rangle \geq \langle p, \sum_{i \in I} e_i \rangle$, we must also have $\langle p, \lambda x_i \rangle \geq \langle p, e_i \rangle$. Dividing through by λ gives $\langle p, x_i \rangle \geq \langle \frac{1}{\lambda}p, e_i \rangle > \langle p, e_i \rangle$ where the strict inequality holds since e_i is nonnegative and nonzero while p is positive.

In addition to the weakening of preference monotonicity, we also dispense with the typical assumption required by the existing second welfare theorem that every consumer has a positive endowment for every type of good (i.e., $e_i \gg 0, \forall i \in I$). The positive endowment assumption can be quite restrictive, excluding many realistic situations. Relaxing the same assumption was an important motivation behind Arrow's generalization of the first welfare theorem.³⁰ At the same time, the theorem assumes PLC utility functions. This assumption guarantees that the "upper contour set" of the target allocation—or the set of goods weakly preferred to $(e_i)_{i\in I}$ —is a polyhedron. Meanwhile, preference monotonicity and Pareto-optimality of $(u_i(e_i))_{i\in I}$ ensure that the vector \bar{e} is a (minimal) face of this set. Invoking Proposition D.2, \bar{e} is then exposed by a positive normal (or price vector) that supports $(e_i)_{i\in I}$ as a competitive equilibrium allocation.

³⁰ "While listening to a talk about housing by Franko (sic) Modigliani, Arrow realized that most people consume nothing of most goods (for example, living in just one particular kind of house), and thus that the prevailing efficiency proofs assumed away all the realistic cases," according to Geanakoplos in https://www.econometricsociety.org/sites/default/files/inmemoriam/arrow_geanakoplos.pdf.